



Abstract Linear Algebra - Part 6

subset of $\mathcal{F}(\mathbb{R})$ given by:

$$\cos: \mathbb{R} \rightarrow \mathbb{R} \rightsquigarrow \text{graph of } \cos(x)$$

$$\sin: \mathbb{R} \rightarrow \mathbb{R} \rightsquigarrow \text{graph of } \sin(x)$$

$$\exp: \mathbb{R} \rightarrow \mathbb{R} \rightsquigarrow \text{graph of } \exp(x)$$

$$\mathcal{U} := \text{span}(\cos, \sin, \exp)$$

Question: Is (\cos, \sin, \exp) a basis of \mathcal{U} ?

generating ✓

linearly independent ?

We have to check: $\alpha_1 \cdot \cos + \alpha_2 \cdot \sin + \alpha_3 \cdot \exp = 0 \Rightarrow \alpha_j = 0$ for all j

means:

zero vector in $\mathcal{F}(\mathbb{R})$

$$\alpha_1 \cdot \cos(x) + \alpha_2 \cdot \sin(x) + \alpha_3 \cdot \exp(x) = 0(x)$$

$$\hookrightarrow 0: \mathbb{R} \rightarrow \mathbb{R}$$

$$x \mapsto 0$$

for all $x \in \mathbb{R}$

$$\Rightarrow \begin{cases} \alpha_1 \cdot \cos(0) + \alpha_2 \cdot \sin(0) + \alpha_3 \cdot \exp(0) = 0 \\ \alpha_1 \cdot \cos(\frac{\pi}{2}) + \alpha_2 \cdot \sin(\frac{\pi}{2}) + \alpha_3 \cdot \exp(\frac{\pi}{2}) = 0 \\ \alpha_1 \cdot \cos(-2\pi \cdot 500) + \alpha_2 \cdot \sin(-2\pi \cdot 500) + \alpha_3 \cdot \exp(-2\pi \cdot 500) = 0 \end{cases}$$

$$\Rightarrow \begin{cases} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & e^{\pi/2} \\ 1 & 0 & e^{-1000\pi} \end{pmatrix} \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \end{cases} \text{ system of linear equations}$$

$$\text{since } \det \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & e^{\pi/2} \\ 1 & 0 & e^{-1000\pi} \end{pmatrix} = e^{-1000\pi} + 0 + 0 - 1 - 0 - 0 < 0,$$

the system of linear equations is uniquely solvable.

$$\Rightarrow \alpha_1 = \alpha_2 = \alpha_3 = 0 \Rightarrow \text{basis } \mathcal{B} = (\cos, \sin, \exp) \text{ of } \mathcal{U}$$

Basis isomorphism: $\Phi_{\mathcal{B}}: \mathcal{U} \rightarrow \mathbb{R}^3$,

$$\text{defined by } \Phi_{\mathcal{B}}(\cos) = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \Phi_{\mathcal{B}}(\sin) = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \Phi_{\mathcal{B}}(\exp) = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

What about $v: \mathbb{R} \rightarrow \mathbb{R}$, $v(x) = 7 \cos(x) + 2 \exp(x)$

$$\Phi_{\mathcal{B}}(v) = \begin{pmatrix} 7 \\ 0 \\ 2 \end{pmatrix}$$

\mathcal{U} is completely represented by \mathbb{R}^3