


## Abstract Linear Algebra - Part 15

Example:  $\mathbb{R}^2$  with standard inner product.



$$r = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad \hat{r} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

(unit vector)

$$x = \begin{pmatrix} 5 \\ 4 \end{pmatrix}$$


$$p = \frac{\langle r, x \rangle}{\langle r, r \rangle} \cdot r = \langle \hat{r}, x \rangle \cdot \hat{r}$$

(orthogonal projection:  
 $\hat{r} \langle \hat{r}, \cdot \rangle$ )

$$\Rightarrow p = \frac{1}{\sqrt{2}} (5+4) \cdot \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 9/2 \\ 9/2 \end{pmatrix}$$

$$\Rightarrow n = x - p = \begin{pmatrix} 1/2 \\ -1/2 \end{pmatrix}$$

Generalization:



$$x = p + n, \quad p \in U, \quad n \in U^\perp$$

Important fact:  $U \cap U^\perp = \{0\}$  for every subspace  $U \subseteq V$

Proposition:  $V$   $\mathbb{F}$ -vector space, inner product  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ .

Let  $U \subseteq V$  be a  $k$ -dimensional subspace,  $\mathcal{B} = (b_1, b_2, \dots, b_k)$  basis of  $U$ .

Then for  $y \in V$ :  $y \perp U$  for all  $u \in U$

$\Leftrightarrow$

$$y \perp b_j \text{ for all } j \in \{1, 2, \dots, k\}$$

Proof:  $(\Rightarrow) \checkmark$   $(\Leftarrow)$  We assume:  $\langle y, b_j \rangle = 0$  for all  $j \in \{1, 2, \dots, k\}$


$$\Rightarrow \sum_{j=1}^k \lambda_j \langle y, b_j \rangle = 0$$

$$\Rightarrow \left\langle y, \sum_{j=1}^k \lambda_j b_j \right\rangle = 0 \xRightarrow{\mathcal{B} \text{ basis}} y \perp u \text{ for all } u \in U$$

Orthogonal projection onto a subspace:

$V$   $\mathbb{F}$ -vector space, inner product  $\langle \cdot, \cdot \rangle$ ,

$U \subseteq V$   $k$ -dimensional subspace.



For  $x \in V$  and a decomposition  $x = p + n$  with  $p \in U$ ,  $n \in U^\perp$ ,

we call:

$p$  orthogonal projection of  $x$  onto  $U$

$n$  normal component of  $x$  with respect to  $U$