




Abstract Linear Algebra - Part 22



Recall: $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ linear \iff matrix $A \in \mathbb{R}^{m \times n}$

Definition: Let V, W be two \mathbb{F} -vector spaces. (same \mathbb{F} for both)

A map $f: V \rightarrow W$ is called linear if:

$$(1) f(u+v) = f(u) + f(v)$$

vector addition in V


vector addition in W

$$(2) f(\lambda \cdot u) = \lambda \cdot f(u)$$

scalar multiplication in V

scalar multiplication in W

for all $u, v \in V, \lambda \in \mathbb{F}$.



Remember: $f(0_V) = f(0 \cdot u) \stackrel{(2)}{=} 0 \cdot f(u) = 0_W$

Example: (a) $V = \mathbb{F}^3$, $W = \mathbb{F}$, $a \in V$.

$f(u) := \langle a, u \rangle_{\text{standard}}$ is a linear map.

$\equiv a^* u$ (matrix multiplication)

$$\begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}^* = (\overline{a_1} \ \overline{a_2} \ \overline{a_3})$$

(transpose + complex conjugation)

(b) $V = \mathcal{P}_3(\mathbb{R})$, $W = \mathcal{P}_2(\mathbb{R})$

$$l: V \rightarrow W$$

$$p \mapsto p'$$

$$l(x \mapsto x^2) = x \mapsto 2x$$

is a linear map!

$$l(p+q) = (p+q)' = p' + q' = l(p) + l(q)$$

$$l(\lambda p) = (\lambda p)' = \lambda p' = \lambda l(p)$$