




Abstract Linear Algebra - Part 25

$l: V \rightarrow W$  linear:  $l(x+y) = l(x) + l(y)$   
 $l(\lambda \cdot x) = \lambda \cdot l(x)$



$B = (b_1, b_2, \dots, b_n)$  basis of  $V$   
 $\Rightarrow x \in V$  can be written as  $x = \alpha_1 b_1 + \dots + \alpha_n b_n$

Hence:  $l(x) = l(\alpha_1 b_1 + \dots + \alpha_n b_n)$   
 $= \alpha_1 l(b_1) + \alpha_2 l(b_2) + \dots + \alpha_n l(b_n)$   
 If know these, we know  $l$


Example:  $V = P_3(\mathbb{R})$ ,  $W = P_2(\mathbb{R})$ ,  $l: V \rightarrow W$   
 $p \mapsto p'$   
 is a linear map!

basis:  $B = (b_1, b_2, b_3, b_4) = (m_0, m_1, m_2, m_3)$   
 with  $m_0: x \mapsto 1$ ,  $m_k: x \mapsto x^k$

$l(m_0) = 0 \leftarrow$  zero vector:  $x \mapsto 0$

$l(m_k) = k \cdot m_{k-1}$ ,  $k \in \{1, 2, 3\}$

Result:



can be represented by a matrix  
 $f(x) = Ax$   
 $A \in \mathbb{F}^{m \times n}$

First column of  $A$ :  $f(e_1) = (\Phi_C \circ l \circ \Phi_B^{-1})(e_1) = \Phi_C(l(b_1))$

Matrix representation: For a linear map  $l: V \rightarrow W$ ,

$l_{C \leftarrow B} := \begin{pmatrix} | & | & & | \\ \Phi_C(l(b_1)) & \Phi_C(l(b_2)) & \dots & \Phi_C(l(b_n)) \\ | & | & & | \end{pmatrix} \in \mathbb{F}^{m \times n}$

is called the matrix representation of  $l$  with respect to  $B$  and  $C$ .

Example (from before)  $V = P_3(\mathbb{R})$  basis:  $B = (b_1, b_2, b_3, b_4) = (m_0, m_1, m_2, m_3)$   
 with  $m_0: x \mapsto 1$ ,  $m_k: x \mapsto x^k$

$W = P_2(\mathbb{R})$  basis:  $C = (c_1, c_2, c_3) = (m_0, m_1, m_2)$   
 $l: V \rightarrow W$   
 $p \mapsto p'$

is a linear map:

$\Phi_C(l(b_1)) = \Phi_C(l(m_0)) = \Phi_C(0) = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \in \mathbb{F}^3$

$\Phi_C(l(b_2)) = \Phi_C(l(m_1)) = \Phi_C(m_0) = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \in \mathbb{F}^3$

$\vdots$

$\Rightarrow l_{C \leftarrow B} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix}$  matrix representation of  $l$