



## Abstract Linear Algebra - Part 29

Equivalence relation:  $A, B \in \mathbb{F}^{m \times n}$ ,  $A \sim B$   $\leftarrow$  they both represent the same linear map  $\ell: V \rightarrow W$

$\leftarrow$  there are invertible matrices  $S, T$  with  $B = S A T$ .

kernel and range?

$$\text{Ker}(B) = \text{Ker}(SAT) = \{x \in \mathbb{F}^n \mid \underbrace{ATx}_{\in \text{Ker}(A)} = 0\} = T^{-1} \text{Ker}(A)$$

$$\begin{aligned} \text{Ran}(B) &= \text{Ran}(SAT) = \{SATx \mid x \in \mathbb{F}^n\} \\ &= \{SA\tilde{x} \mid \tilde{x} \in \mathbb{F}^n\} = S \text{Ran}(A) \end{aligned}$$

Result:  $A \sim B \implies$

$$\begin{array}{ccc} \text{rank}(A) & = & \text{rank}(B) \\ + & & + \\ \text{nullity}(A) & = & \text{nullity}(B) \\ \parallel & & \parallel \\ n & & n \end{array}$$

Proposition: For  $A, B \in \mathbb{F}^{m \times n}$ , we have:

$$A \sim B \iff \text{rank}(A) = \text{rank}(B)$$

Proof:

$$A \sim \left( \begin{array}{c|ccc} \square & & & \\ \hline & \square & & \\ & & \square & \\ & & & \square \end{array} \right) \xrightarrow{\text{Gaussian elimination}} \left( \begin{array}{c|ccc} \square & 0 & 0 & \dots \\ \hline & \square & & \\ & & \square & \\ & & & \square \end{array} \right) \xrightarrow{\text{backwards substitution}} \left( \begin{array}{c|ccc} \square & & & \\ \hline & \square & & \\ & & \square & \\ & & & \square \end{array} \right) \xrightarrow{\text{column/row exchanges}} \left( \begin{array}{c|ccc} \square & & & \\ \hline & \square & & \\ & & \square & \\ & & & \square \end{array} \right)$$

$$\begin{aligned} \implies A &\sim \begin{pmatrix} \mathbb{1}_r & 0 \\ 0 & 0 \end{pmatrix} & \text{with } r &= \text{rank}(A) \\ B &\sim \begin{pmatrix} \mathbb{1}_r & 0 \\ 0 & 0 \end{pmatrix} & r &= \text{rank}(B) \end{aligned}$$

□