



## Algebra - Part 14

Recall: Symmetric group  $S_3$



Definition: Let  $G$  be a group and  $S \subseteq G$  be a subset.

$$\langle S \rangle := \bigcap_{\substack{U \subseteq G \text{ subgroup} \\ \text{with } S \subseteq U}} U$$



We say:  $S$  generates the subgroup  $\langle S \rangle$ .

Proposition: Intersection of subgroups is also a subgroup.

Proof: Assume:  $G$  group,  $U_j \subseteq G$  subgroups for all  $j \in J$ ,  $\tilde{U} := \bigcap_{j \in J} U_j$ .

Obvious:  $e \in \tilde{U}$  ✓

Take  $a, b \in \tilde{U} \implies a, b \in U_j$  for all  $j \in J$

$\implies ab \in U_j$  and  $a^{-1} \in U_j$  for all  $j \in J$

$\implies ab \in \tilde{U}$  and  $a^{-1} \in \tilde{U}$  □

Fact: If  $S \neq \emptyset$  and  $S^{-1} := \{s^{-1} \mid s \in S\}$ , then:

$$\langle S \rangle = \{a_1 a_2 \cdots a_n \in G \mid n \in \mathbb{N}, a_1, \dots, a_n \in S \cup S^{-1}\}$$

Example: Symmetric group  $S_3$ :  $S = \{a, b\}$ ,  $S^{-1} = \{a^2, b\}$

$\leadsto ab, ba, bb = e, \dots$  just six elements

$$S_3 = \langle a, b \rangle$$

Definition: A group  $G$  is called cyclic if there is element  $g \in G$

such that  $\langle g \rangle = G$ .

In other words:  $G = \{g^k \mid k \in \mathbb{Z}\}$  with  $g^0 :=$  identity element in  $G$