

## Distributions - Part 3

### Convergence for test functions

$$\mathcal{D}(\mathbb{R}^n) = C_c^\infty(\mathbb{R}^n) \quad (\text{space of test functions})$$

Notions of convergence:

• If one has a norm  $\|\cdot\|$ :

$$f_n \rightarrow f \quad :\Leftrightarrow \quad \|f_n - f\| \xrightarrow{n \rightarrow \infty} 0$$

• If one has a metric  $d(\cdot, \cdot)$ :


$$f_n \rightarrow f \quad :\Leftrightarrow \quad d(f_n, f) \xrightarrow{n \rightarrow \infty} 0$$

$\Rightarrow$  Both notions are not sufficient here  $\rightsquigarrow$  very strong notion needed

Definition: For  $(\varphi_k)_{k \in \mathbb{N}} \subset \mathcal{D}(\mathbb{R}^n)$ ,  $\varphi \in \mathcal{D}(\mathbb{R}^n)$ , we write


$$\varphi_k \xrightarrow{\mathcal{D}} \varphi \quad \text{if}$$

(a) There is a bounded set  $M$ ,  
such that outside of it:  
 $\varphi_k = 0$  for all  $k$ .



(b) Uniform convergence  $\varphi_k \xrightarrow{\text{unif.}} \varphi$   
and for all multi-indices  $\alpha$

$$D^\alpha \varphi_k \xrightarrow{\text{unif.}} D^\alpha \varphi$$



Use the supremum norm  $\|f\|_\infty := \sup \{ |f(x)| \mid x \in \mathbb{R}^n \}$ .

$$\psi_k \xrightarrow{\mathcal{D}} \psi \quad \Leftrightarrow \quad \begin{aligned} & \text{(a) } \exists C \subseteq \mathbb{R}^n \text{ compact such that} \\ & \quad \text{supp}(\psi_1), \text{supp}(\psi_2), \dots \subseteq C \\ & \text{(b) } \forall \alpha \text{ multi-index we have} \\ & \quad \|D^\alpha \psi_k - D^\alpha \psi\|_\infty \xrightarrow{k \rightarrow \infty} 0 \end{aligned}$$