

The Bright Side of Mathematics


Fourier Transform - Part 14

$$\|f - \mathcal{F}_n(f)\|_{L^2} \xrightarrow{n \rightarrow \infty} 0$$

not a pointwise convergence!

~> We can get uniform convergence for special functions

Example: continuous and piecewise C^1 -function




Supremum norm: $\|f\|_\infty := \sup_{x \in [-\pi, \pi]} |f(x)|$

$$\Rightarrow \int_{-\pi}^{\pi} |f(x)|^2 dx \leq \int_{-\pi}^{\pi} \|f\|_\infty^2 dx = 2\pi \cdot \|f\|_\infty^2$$

$$\Rightarrow \|f\|_{L^2} \leq \|f\|_\infty$$

Theorem: $f: \mathbb{R} \rightarrow \mathbb{C}$ 2π -periodic continuous function.



Assume there are finitely many points (a_1, a_2, \dots, a_m)

inside the interval $[-\pi, \pi]$ such that:

$$f|_{[a_j, a_{j+1}]} \in C^1 \quad \text{for all } j \in \{0, 1, \dots, m\}$$

Then: $\|f - \mathcal{F}_n(f)\|_\infty \xrightarrow{n \rightarrow \infty} 0$

$$\left(\begin{array}{l} \mathcal{F}_n(f) = \sum_{k=-n}^n e_k \langle e_k, f \rangle \\ e_k: x \mapsto e^{ikx} \\ \langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \overline{f(x)} \cdot g(x) dx \end{array} \right)$$