


The Bright Side of Mathematics

The following pages cover the whole Fourier Transform course of the Bright Side of Mathematics. Please note that the creator lives from generous supporters and would be very happy about a donation. See more here: <https://tbsom.de/support>

Have fun learning mathematics!



Fourier Series


Exercises 1

Exercise 1. Compute the Fourier series of $f(x) = |\sin(x)|$.

$$f(x) \approx \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos(k\omega x) + b_k \sin(k\omega x)) \quad \omega = \frac{2\pi}{T}$$

$$a_k = \frac{2}{T} \int_0^T f(x) \cos(k\omega x) dx, \quad k \geq 0$$

$$b_k = \frac{2}{T} \int_0^T f(x) \sin(k\omega x) dx, \quad k \geq 1$$



→ even: $b_k = 0$

$$T = \pi \quad \omega = \frac{2\pi}{\pi} = 2$$

$$\frac{a_0}{2} = \frac{1}{\pi} \int_0^{\pi} \sin(x) dx = \frac{1}{\pi} (-\cos^1 \pi + \cos^1 0) = \frac{2}{\pi}$$

$$a_k = \frac{2}{\pi} \int_0^{\pi} \sin(x) \cos(2kx) dx$$

$$\int \sin(x) \cos(2kx) dx = -\cos(x) \cos(2kx) - 2k \int \cos(x) \sin(2kx) dx$$

$$f'(x) = \sin(x) \quad g(x) = \cos(2kx)$$

$$f'(x) = \cos x \quad g(x) = \sin(2kx)$$

$$f(x) = -\cos(x) \quad g'(x) = -\sin(2kx) 2k$$

$$f(x) = \sin x \quad g'(x) = \cos(2kx) 2k$$

$$\int \sin(x) \cos(2kx) dx = -\cos(x) \cos(2kx) - 2k \left(\sin(x) \sin(2kx) - 2k \int \sin x \cos(2kx) dx \right)$$


$$(1 - 4k^2) \int_0^{\pi} \sin(x) \cos(2kx) dx = \left(-\cos(x) \cos(2kx) - 2k \sin(x) \sin(2kx) \right) \Big|_0^{\pi}$$

$$\int_0^{\pi} \sin(x) \cos(2kx) dx = \frac{1}{1-4k^2} \left(\underbrace{-(-1)(1)}_1 + \underbrace{-(-1)(1)}_1 \right) = \frac{2}{1-4k^2}$$

$$a_k = \frac{2}{\pi} \cdot \frac{2}{1-4k^2} \quad k \geq 1, \quad \frac{a_0}{2} = \frac{2}{\pi}, \quad b_k = 0$$


$$f(x) \approx \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos(k\omega x) + b_k \sin(k\omega x))$$

$$|\sin(x)| \approx \frac{2}{\pi} + \sum_{k=1}^{\infty} \frac{4}{\pi(1-4k^2)} \cos(2kx)$$




Fourier Transform - Part 2

Idea of Fourier series:






The function is 2-periodic: $f(x+2) = f(x)$ for all $x \in \mathbb{R}$



$$\mathcal{F}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{R}) = \{ f: \mathbb{R} \rightarrow \mathbb{R} \mid f(x+2\pi) = f(x) \text{ for all } x \in \mathbb{R} \}$$

↳ real vector space

Example: • constant function $f(x) = 5$



$$x \mapsto \sin(x)$$

$$x \mapsto \sin(2x)$$

Proposition: $U \subseteq \mathcal{F}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{R})$ given by

$$U := \left\{ \begin{array}{l} x \mapsto \sin(x), x \mapsto \sin(2x), x \mapsto \sin(3x), \dots, \\ x \mapsto 1, x \mapsto \cos(x), x \mapsto \cos(2x), x \mapsto \cos(3x), \dots \end{array} \right\}$$

odd functions ↙
↖ even functions


is linearly independent.

Definition: A linear combination $f \in \text{Span}(U)$, $f: \mathbb{R} \rightarrow \mathbb{R}$, is called
(real) trigonometric polynomial:

$$f(x) = a_0 + \sum_{k=1}^n a_k \cdot \cos(k \cdot x) + \sum_{k=1}^n b_k \cdot \sin(k \cdot x), \quad a_i, b_i \in \mathbb{R}$$

For $\mathcal{F}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$, we have a (complex) trigonometric polynomial:

$$f(x) = \sum_{k=-n}^n c_k \cdot \exp(i \cdot k \cdot x), \quad c_k \in \mathbb{C}$$



Fourier Transform - Part 3

In $\mathcal{F}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{R})$, we have (real) trigonometric polynomials:

$$f(x) = a_0 + \sum_{k=1}^n a_k \cdot \cos(k \cdot x) + \sum_{k=1}^n b_k \cdot \sin(k \cdot x) \quad , \quad a_i, b_i \in \mathbb{R}$$

Subspace: $\mathcal{P}_{2\pi\text{-per}} := \text{span} \left(\begin{array}{l} x \mapsto 1, \quad x \mapsto \cos(x), \quad x \mapsto \cos(2x), \quad x \mapsto \cos(3x), \dots, \\ x \mapsto \sin(x), \quad x \mapsto \sin(2x), \quad x \mapsto \sin(3x), \dots \end{array} \right)$

basis!

Definition: For $f, g \in \mathcal{P}_{2\pi\text{-per}}$, we define an inner product:

$$\langle f, g \rangle := \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) g(x) dx$$

Example: $\langle x \mapsto 1, x \mapsto 1 \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} 1 dx = 1$

$$\begin{aligned} \langle x \mapsto \cos(x), x \mapsto \sin(x) \rangle &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \cos(x) \sin(x) dx \\ &= \frac{1}{2\pi} \left(\frac{1}{2} (\sin(x))^2 \Big|_{-\pi}^{\pi} \right) = 0 \end{aligned}$$

$$\langle x \mapsto \cos(k \cdot x), x \mapsto \sin(m \cdot x) \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \underbrace{\cos(k \cdot x) \sin(m \cdot x)}_{\text{odd function}} dx = 0$$


$$\langle x \mapsto 1, x \mapsto \cos(k \cdot x) \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \cos(k \cdot x) dx = \frac{1}{2\pi} \frac{1}{k} \sin(k \cdot x) \Big|_{-\pi}^{\pi} = 0$$

$$\langle x \mapsto 1, x \mapsto \sin(m \cdot x) \rangle = 0$$

$$\langle x \mapsto \cos(k \cdot x), x \mapsto \cos(m \cdot x) \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \cos(k \cdot x) \cos(m \cdot x) dx$$

$$= 0 \quad \text{if } k \neq m$$

$$\text{Use: } \cos(x) = \frac{1}{2} (e^{ix} + e^{-ix})$$



$$\text{Then: } \int_{-\pi}^{\pi} \cos(k \cdot x) \cos(m \cdot x) dx = \frac{1}{4} \int_{-\pi}^{\pi} \left(e^{i(k+m)x} + e^{-i(k+m)x} + e^{i(k-m)x} + e^{-i(k-m)x} \right) dx$$

$$\stackrel{k \neq m}{=} \frac{1}{4} \left(\frac{1}{i(k+m)} e^{i(k+m)x} + \frac{1}{-i(k+m)} e^{-i(k+m)x} + \frac{1}{i(k-m)} e^{i(k-m)x} + \frac{1}{-i(k-m)} e^{-i(k-m)x} \right) \Big|_{-\pi}^{\pi}$$

$$\text{Use: } \sin(x) = \frac{1}{2i} (e^{ix} - e^{-ix})$$

$$= \frac{1}{2} \left(\frac{1}{k+m} \sin((k+m) \cdot x) + \frac{1}{k-m} \sin((k-m) \cdot x) \right) \Big|_{-\pi}^{\pi} = 0$$


$$\text{And similarly: } \int_{-\pi}^{\pi} \sin(k \cdot x) \sin(m \cdot x) dx \stackrel{k \neq m}{=} 0$$

Result: $\mathcal{B} = \left(x \mapsto 1, x \mapsto \cos(x), x \mapsto \cos(2x), x \mapsto \cos(3x), \dots, \right.$
 $\left. x \mapsto \sin(x), x \mapsto \sin(2x), x \mapsto \sin(3x), \dots \right)$

satisfies $\langle f, g \rangle = 0 \quad f \neq g, f, g \in \mathcal{B}$

$\leadsto \mathcal{B}$ orthogonal basis (OB)

\leadsto make to orthonormal basis (ONB)



Fourier Transform - Part 4

We already know: we have an orthogonal basis (OB)

$$\mathcal{B} = \left(\begin{array}{l} x \mapsto 1, \quad x \mapsto \cos(x), \quad x \mapsto \cos(2x), \quad x \mapsto \cos(3x), \dots, \\ x \mapsto \sin(x), \quad x \mapsto \sin(2x), \quad x \mapsto \sin(3x), \dots \end{array} \right)$$

for $\mathcal{P}_{2\pi\text{-per}}$ with inner product $\langle f, g \rangle_1 := \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)g(x) dx$

Normalize:

$$\langle x \mapsto \sin(kx), x \mapsto \sin(kx) \rangle_1 = \frac{1}{2\pi} \int_{-\pi}^{\pi} (\sin(kx))^2 dx \quad \begin{array}{c} \uparrow \\ \square \\ \rightarrow \end{array}$$

$$\int_{-\pi}^{\pi} (\sin(kx))^2 dx = \int_{-\pi}^{\pi} \underbrace{\sin(kx)}_u \underbrace{\sin(kx)}_{v'} dx = \left. \sin(kx) \left(-\frac{1}{k}\right) \cos(kx) \right|_{-\pi}^{\pi} - \int_{-\pi}^{\pi} k \cos(kx) \left(-\frac{1}{k}\right) \cos(kx) dx$$

integration by parts: $u' = k \cos(kx)$
 $v = -\frac{1}{k} \cos(kx)$

$$= \int_{-\pi}^{\pi} \underbrace{(\cos(kx))^2}_{1 - (\sin(kx))^2} dx$$

$$\Rightarrow 2 \cdot \int_{-\pi}^{\pi} (\sin(kx))^2 dx = \int_{-\pi}^{\pi} 1 dx = 2\pi$$

$$\langle x \mapsto \sin(kx), x \mapsto \sin(kx) \rangle_1 = \frac{1}{2} \rightsquigarrow \text{length} = \frac{1}{\sqrt{2}}$$

Hence: $x \mapsto \sqrt{2} \cdot \sin(kx)$ has norm 1

Proposition: (1) $\mathcal{B} = \left(\begin{array}{l} x \mapsto 1, \quad x \mapsto \sqrt{2} \cos(x), \quad x \mapsto \sqrt{2} \cos(2x), \quad x \mapsto \sqrt{2} \cos(3x), \dots, \\ x \mapsto \sqrt{2} \sin(x), \quad x \mapsto \sqrt{2} \sin(2x), \quad x \mapsto \sqrt{2} \sin(3x), \dots \end{array} \right)$

is an ONB w.r.t. the inner product: $\langle f, g \rangle_1 := \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)g(x) dx$

$$(2) \quad \mathcal{B} = \left(x \mapsto \frac{1}{\sqrt{2\pi}}, x \mapsto \frac{1}{\sqrt{\pi}} \cos(x), x \mapsto \frac{1}{\sqrt{\pi}} \cos(2x), x \mapsto \frac{1}{\sqrt{\pi}} \cos(3x), \dots, \right. \\ \left. x \mapsto \frac{1}{\sqrt{\pi}} \sin(x), x \mapsto \frac{1}{\sqrt{\pi}} \sin(2x), x \mapsto \frac{1}{\sqrt{\pi}} \sin(3x), \dots \right)$$

is an ONB w.r.t. the inner product: $\langle f, g \rangle_2 := \int_{-\pi}^{\pi} f(x)g(x) dx$

$$(3) \quad \mathcal{B} = \left(x \mapsto \frac{1}{\sqrt{2}}, x \mapsto \cos(x), x \mapsto \cos(2x), x \mapsto \cos(3x), \dots, \right. \\ \left. x \mapsto \sin(x), x \mapsto \sin(2x), x \mapsto \sin(3x), \dots \right)$$

is an ONB w.r.t. the inner product: $\langle f, g \rangle_3 := \frac{1}{\sqrt{2}} \int_{-\pi}^{\pi} f(x)g(x) dx$

For trigonometric polynomials:

$$f(x) = \tilde{a}_0 \frac{1}{\sqrt{2}} + \sum_{k=1}^n a_k \cos(k \cdot x) + \sum_{k=1}^n b_k \sin(k \cdot x), \quad a_i, b_i \in \mathbb{R}$$

Fourier coefficients w.r.t. ONB in (3)


$$a_k = \langle x \mapsto \cos(k \cdot x), f \rangle_3, \quad \tilde{a}_0 = \langle x \mapsto \frac{1}{\sqrt{2}}, f \rangle_3$$

$$b_k = \langle x \mapsto \sin(k \cdot x), f \rangle_3$$

Approximation of periodic functions?

$$g : \mathbb{R} \rightarrow \mathbb{R}$$

2π -periodic + "integrable"




trigonometric polynomials with basis:

$$\mathcal{B} = (h_1, h_2, \dots, h_N)$$

ONB!

$$\text{orthogonal projection} = \sum_{k=1}^N h_k \langle h_k, g \rangle$$



Fourier Transform - Part 5

$$\mathcal{F}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) = \left\{ f: \mathbb{R} \rightarrow \mathbb{C} \mid f(x+2\pi) = f(x) \text{ for all } x \in \mathbb{R} \right\}$$

$$\mathcal{P}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) := \text{span} \left(x \mapsto \frac{1}{\sqrt{2}}, x \mapsto \cos(x), x \mapsto \cos(2x), x \mapsto \cos(3x), \dots, \right. \\ \left. x \mapsto \sin(x), x \mapsto \sin(2x), x \mapsto \sin(3x), \dots \right)$$

$$\hookrightarrow \text{inner product } \langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \overline{f(x)} g(x) dx$$


Let's take integrable functions:

$$\mathcal{L}_{2\pi\text{-per}}^1(\mathbb{R}, \mathbb{C}) = \left\{ f \in \mathcal{F}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \mid \underbrace{\int_{-\pi}^{\pi} |f(x)| dx}_{f \text{ integrable with respect to Lebesgue measure on } [-\pi, \pi]} < \infty \right\}$$

\hookrightarrow complex vector space

norm? $\|f\|_1 := \int_{-\pi}^{\pi} |f(x)| dx$ problem:

\hookrightarrow not a norm on $\mathcal{L}_{2\pi\text{-per}}^1(\mathbb{R}, \mathbb{C})$



solution: equivalence relation $f \sim g : \Leftrightarrow \|f-g\|_1 = 0$

set of all equivalence classes: $\mathcal{L}_{2\pi\text{-per}}^1(\mathbb{R}, \mathbb{C}) := \mathcal{L}_{2\pi\text{-per}}^1(\mathbb{R}, \mathbb{C}) / \sim$

\hookrightarrow complex vector space

$$\|[f]\|_1 := \|f\|_1$$

\hookrightarrow norm!

identify: $\mathcal{L}_{2\pi\text{-per}}^1(\mathbb{R}, \mathbb{C}) \supseteq \mathcal{P}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$

Let's take square-integrable functions:


$$\mathcal{L}_{2\pi\text{-per}}^2(\mathbb{R}, \mathbb{C}) = \left\{ f \in \mathcal{F}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \mid \int_{-\pi}^{\pi} |f(x)|^2 dx < \infty \right\}$$

norm? $\|f\|_2 := \sqrt{\int_{-\pi}^{\pi} |f(x)|^2 dx}$

solution: equivalence relation $f \sim g : \Leftrightarrow \|f - g\|_2 = 0$

set of all equivalence classes: $\mathcal{L}_{2\pi\text{-per}}^2(\mathbb{R}, \mathbb{C}) := \mathcal{L}_{2\pi\text{-per}}^2(\mathbb{R}, \mathbb{C}) / \sim$

↳ complex vector space with inner product




Fourier Transform - Part 6

We know: $L^1_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \supseteq L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \supseteq \mathcal{P}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$

inner product: $\langle f, g \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} \overline{f(x)} \cdot g(x) dx$

Orthogonality: $\mathcal{B}_n = \left(x \mapsto \frac{1}{\sqrt{2}}, x \mapsto \cos(x), x \mapsto \cos(2x), \dots, x \mapsto \cos(nx) \right.$
 $\left. x \mapsto \sin(x), x \mapsto \sin(2x), x \mapsto \sin(3x), \dots, x \mapsto \sin(nx) \right)$
ONS in $L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ for every $n \in \mathbb{N}$



U_n finite-dimensional subspace spanned by \mathcal{B}_n

write: $\mathcal{B}_n = (h_1, h_2, \dots, h_N)$, $N = 2n + 1$

orthogonal projection of f onto U_n :

$$\mathcal{F}_n(f) = \sum_{k=1}^N h_k \underbrace{\langle h_k, f \rangle}_{\text{Fourier coefficients}}$$

Fourier coefficients

Definition:

$$\mathcal{F}_n(f)(x) = \tilde{a}_0 \frac{1}{\sqrt{2}} + \sum_{k=1}^n a_k \cos(k \cdot x) + \sum_{k=1}^n b_k \sin(k \cdot x)$$

$$\text{with } \tilde{a}_0 = \left\langle x \mapsto \frac{1}{\sqrt{2}}, f \right\rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} \frac{1}{\sqrt{2}} f(x) dx$$


$$a_k = \left\langle x \mapsto \cos(k \cdot x), f \right\rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} \cos(k \cdot x) f(x) dx$$

$$b_k = \left\langle x \mapsto \sin(k \cdot x), f \right\rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} \sin(k \cdot x) f(x) dx$$

The map $h \mapsto \mathcal{F}_n(f)(x)$ (with $x \in \mathbb{R}$)

is called the Fourier series of $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ (can be extended to $f \in L^1_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$)

Example: $f: \mathbb{R} \rightarrow \mathbb{C}$, $f(x) = \begin{cases} 1, & x \in (-\pi, 0) \\ 0, & x \in [0, \pi] \end{cases}$




$$\tilde{a}_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} \frac{1}{\sqrt{2}} f(x) dx = \frac{1}{\pi} \int_{-\pi}^0 \frac{1}{\sqrt{2}} dx = \frac{1}{\sqrt{2}}$$


$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} \cos(k \cdot x) f(x) dx = \frac{1}{\pi} \int_{-\pi}^0 \cos(k \cdot x) dx = 0$$

$$b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} \sin(k \cdot x) f(x) dx = \frac{1}{\pi} \int_{-\pi}^0 \sin(k \cdot x) dx = \frac{1}{\pi} \left(-\frac{1}{k} \cos(k \cdot x) \right) \Big|_{-\pi}^0$$


$$= \begin{cases} 0, & k \text{ even} \\ -\frac{2}{\pi k}, & k \text{ odd} \end{cases}$$

Fourier series: $\frac{1}{2} + \frac{-2}{\pi} \sin(x) + \frac{-2}{\pi 3} \cdot \sin(3 \cdot x) + \frac{-2}{\pi 5} \cdot \sin(5 \cdot x) + \dots$





Fourier Transform - Part 7



Euler's formula: $\cos(x) = \frac{1}{2} (e^{ix} + e^{-ix})$

$$\sin(x) = \frac{1}{2i} (e^{ix} - e^{-ix})$$

Example:

$$A \cdot \cos(x) + B \cdot \cos(2x) + C \sin(2x), \quad A, B, C \in \mathbb{C}$$

$$= \frac{A}{2} (e^{ix} + e^{-ix}) + \frac{B}{2} (e^{i2x} + e^{-i2x}) + \frac{C}{2i} (e^{i2x} - e^{-i2x})$$

$$= \frac{A}{2} \cdot e^{ix} + \frac{A}{2} \cdot e^{-ix} + \left(\frac{B}{2} + \frac{C}{2i}\right) e^{i2x} + \left(\frac{B}{2} - \frac{C}{2i}\right) e^{-i2x}$$

complex linear combination!

Remember: In $\mathcal{P}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$:

$$\text{Span} \left(x \mapsto \frac{1}{\sqrt{2}}, x \mapsto \cos(x), x \mapsto \cos(2x), \dots, x \mapsto \cos(nx), \right. \\ \left. x \mapsto \sin(x), x \mapsto \sin(2x), x \mapsto \sin(3x), \dots, x \mapsto \sin(nx) \right)$$

$$= \text{Span} \left(x \mapsto e^{-inx}, \dots, x \mapsto e^{-ix}, x \mapsto e^{i0x}, x \mapsto e^{ix}, \dots, x \mapsto e^{inx} \right)$$

and $\tilde{a}_0 \frac{1}{\sqrt{2}} + \sum_{k=1}^n a_k \cdot \cos(k \cdot x) + \sum_{k=1}^n b_k \cdot \sin(k \cdot x) = \sum_{k=-n}^n c_k e^{ikx}$

$$\text{with } c_k = \begin{cases} \frac{1}{2} \left(a_k + \frac{b_k}{i} \right), & \text{for } k > 0 \\ \tilde{a}_0 \frac{1}{\sqrt{2}} & \text{for } k = 0 \\ \frac{1}{2} \left(a_{-k} - \frac{b_{-k}}{i} \right), & \text{for } k < 0 \end{cases}$$

Result: Take $L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \supseteq \mathcal{P}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$


with inner product: $\langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \overline{f(x)} \cdot g(x) dx$

best factor for exponential functions

ONS: $\mathcal{B}_n = \left(x \mapsto 1, x \mapsto \sqrt{2} \cos(x), x \mapsto \sqrt{2} \cos(2x), x \mapsto \sqrt{2} \cos(3x), \dots, x \mapsto \sqrt{2} \cos(nx), \right. \\ \left. x \mapsto \sqrt{2} \sin(x), x \mapsto \sqrt{2} \sin(2x), x \mapsto \sqrt{2} \sin(3x), \dots, x \mapsto \sqrt{2} \sin(nx) \right)$

ONS: $\mathcal{E}_n = \left(x \mapsto e^{ikx} \right)_{k=-n, \dots, n} = \left(e_k \right)_{k=-n, \dots, n}$

they span the same subspace




For $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$: $\mathcal{F}_n(f) = \sum_{k=-n}^n e_k \underbrace{\langle e_k, f \rangle}_{\text{Fourier coefficients}}$

$\Rightarrow \mathcal{F}_n(f)(x) = \sum_{k=-n}^n c_k e^{ikx}, \quad c_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikx} f(x) dx$

The map $f \mapsto \mathcal{F}_n(f)$ is called the Fourier series of $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$

(with complex coefficients)



Fourier Transform - Part 8


Fourier series: $f \in L^1_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \rightsquigarrow \mathcal{F}_n(f) \in \mathcal{P}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$

trigonometric polynomial

$$\mathcal{F}_n(f) = \sum_{k=-n}^n c_k e^{ikx}$$

$$c_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikx} f(x) dx$$

Geometric picture: For $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \rightsquigarrow \mathcal{F}_n(f) \in \mathcal{P}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$



orthogonal projection

$$\mathcal{F}_n(f) \perp \underbrace{f - \mathcal{F}_n(f)}_{\text{normal component}}$$

normal component

Question: What happens for $n \rightarrow \infty$? $\mathcal{F}_n(f) \xrightarrow{n \rightarrow \infty} f$?

Proposition: $L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ with inner product $\langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \overline{f(x)} \cdot g(x) dx$


and ONS $(\dots, e_{-2}, e_{-1}, e_0, e_1, e_2, \dots)$ given by $e_k: x \mapsto e^{ikx}$.

Then for $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ and $\mathcal{F}_n(f) = \sum_{k=-n}^n e_k \underbrace{\langle e_k, f \rangle}_{c_k}$,

we have:

$$(a) \quad \|f - \mathcal{F}_n(f)\|^2 = \|f\|^2 - \sum_{k=-n}^n |c_k|^2$$

L^2 -norm $\|\cdot\| := \sqrt{\langle \cdot, \cdot \rangle}$




Pythagorean theorem: $\|f\|^2 = \|\mathcal{F}_n(f)\|^2 + \|f - \mathcal{F}_n(f)\|^2$

$$(b) \sum_{k=-n}^n |c_k|^2 \leq \|f\|^2 \quad \text{for all } n \quad (\text{Bessel's inequality})$$

$$\left(\Rightarrow \sum_{k=-\infty}^{\infty} |c_k|^2 \leq \|f\|^2 \quad \text{and} \quad c_k \xrightarrow{k \rightarrow \infty} 0 \right)$$

$$(c) \|f - \mathcal{F}_n(f)\| \xrightarrow{n \rightarrow \infty} 0 \quad \Leftrightarrow \sum_{k=-\infty}^{\infty} |c_k|^2 = \|f\|^2$$

(Parseval's identity)



Fourier Transform - Part 9

$L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ has ONS $(\dots, e_{-2}, e_{-1}, e_0, e_1, e_2, \dots)$ given by $e_k: x \mapsto e^{ikx}$

\rightsquigarrow Fourier series $\mathcal{F}_n(f) = \sum_{k=-n}^n e_k \langle e_k, f \rangle$

Parseval's identity: $\|f\|^2 = \sum_{k=-\infty}^{\infty} |\langle e_k, f \rangle|^2$

$$\Leftrightarrow \|f - \mathcal{F}_n(f)\| \xrightarrow{n \rightarrow \infty} 0$$

means: $f = \mathcal{F}_n(f) + r_n$ with $\|r_n\| \xrightarrow{n \rightarrow \infty} 0$

Consider two functions: $f, g \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$

$\langle f, g \rangle \leftarrow$ formula with Fourier coefficients?

$$f = \mathcal{F}_n(f) + r_n \quad \text{with} \quad \|r_n\| \xrightarrow{n \rightarrow \infty} 0$$

$$g = \mathcal{F}_n(g) + \tilde{r}_n \quad \text{with} \quad \|\tilde{r}_n\| \xrightarrow{n \rightarrow \infty} 0$$

$$\text{We have: } |\langle \mathcal{F}_n(g), r_n \rangle| \leq \|\mathcal{F}_n(g)\| \|r_n\|$$

Cauchy
Schwarz

$$\leq \|g\| \cdot \|r_n\| \xrightarrow{n \rightarrow \infty} 0$$

Bessel's inequality \nearrow

$$\langle f, g \rangle = \langle \mathcal{F}_n(f) + r_n, \mathcal{F}_n(g) + \tilde{r}_n \rangle$$

$$= \langle \mathcal{F}_n(f), \mathcal{F}_n(g) \rangle + \underbrace{\langle r_n, \mathcal{F}_n(g) \rangle + \langle \mathcal{F}_n(f), \tilde{r}_n \rangle}_{(*)} + \langle r_n, \tilde{r}_n \rangle$$

$$= \left\langle \sum_{k=-n}^n e_k \langle e_k, f \rangle, \sum_{l=-n}^n e_l \langle e_l, g \rangle \right\rangle + (*)$$

$$\begin{aligned}
&= \sum_{k=-n}^n \sum_{l=-n}^n \overline{\langle e_k, f \rangle} \langle e_l, g \rangle \underbrace{\langle e_k, e_l \rangle}_{=\delta_{kl}} + (*) \\
&= \sum_{k=-n}^n \langle f, e_k \rangle \langle e_k, g \rangle + (*) \\
&\xrightarrow{h \rightarrow \infty} \sum_{k=-\infty}^{\infty} \langle f, e_k \rangle \langle e_k, g \rangle
\end{aligned}$$

Remember the equivalent statements: $L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ with ONS $(e_k)_{k \in \mathbb{Z}}$

(a) Parseval's identity: $\|f\|^2 = \sum_{k=-\infty}^{\infty} |\langle e_k, f \rangle|^2$

(b) ONS is complete: $\left\| f - \sum_{k=-n}^n e_k \langle e_k, f \rangle \right\| \xrightarrow{h \rightarrow \infty} 0$
 $\left(f = \sum_{k=-\infty}^{\infty} e_k \langle e_k, f \rangle \right)$


(c) ONS gives inner product:


$$\langle f, g \rangle = \sum_{k=-\infty}^{\infty} \langle f, e_k \rangle \langle e_k, g \rangle \quad \left(\sum_{k=-\infty}^{\infty} |e_k\rangle \langle e_k| = \mathbb{1} \right) \quad \text{informal:}$$

(d) ONS is total: $\text{span}\left(\{e_k\}_{k \in \mathbb{Z}}\right)$ is dense in $L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$:

$$\forall f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \quad \forall \epsilon > 0 \quad \exists N \in \mathbb{N}, \lambda_1, \lambda_2, \dots, \lambda_N \in \mathbb{C}:$$


$$\left\| f - \sum_{k=-N}^N \lambda_k e_k \right\| < \epsilon$$





Fourier Transform - Part 10


For proving Parseval's identity \rightsquigarrow step functions



Most important step function:

$$h_a(x) = \begin{cases} 1, & x \in [-\pi, a] \\ 0, & x \in (a, \pi) \end{cases}$$

for every $a \in [-\pi, \pi]$



Fourier series for this example:


$$\begin{aligned} c_k &= \langle e_k, h_a \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikx} h_a(x) dx = \frac{1}{2\pi} \int_{-\pi}^a e^{-ikx} dx \\ &= \begin{cases} \frac{a + \pi}{2\pi}, & k = 0 \\ \frac{1}{2\pi(-ik)} (e^{-ika} - e^{ik\pi}), & k \neq 0 \end{cases} \end{aligned}$$



Visualization:

$$a_k = 2 \cdot \text{Re}(c_k)$$

$$b_k = -2 \cdot \text{Im}(c_k)$$



Show Parseval's identity:

$$\begin{aligned}
 k \neq 0: \quad |c_k|^2 &= \frac{1}{2\tilde{\pi}(-ik)} \left(e^{-ika} - e^{ik\tilde{\pi}} \right) \overline{\frac{1}{2\tilde{\pi}(-ik)} \left(e^{-ika} - e^{ik\tilde{\pi}} \right)} \\
 &= \frac{1}{4\tilde{\pi}^2 k^2} \cdot \left(e^{-ika} - e^{ik\tilde{\pi}} \right) \cdot \left(e^{ika} - e^{-ik\tilde{\pi}} \right) \\
 &= \frac{1}{4\tilde{\pi}^2 k^2} \cdot \left(1 - e^{ik(\tilde{\pi}+a)} - e^{-ik(\tilde{\pi}+a)} + 1 \right) \\
 &= \frac{1}{4\tilde{\pi}^2 k^2} \cdot \left(2 - 2 \cos(k(\tilde{\pi}+a)) \right) = \frac{1}{2\tilde{\pi}^2 k^2} \cdot \left(1 - \cos(k(\tilde{\pi}+a)) \right)
 \end{aligned}$$

$$\Rightarrow \sum_{k=-n}^n |c_k|^2 = \left(\frac{a + \tilde{\pi}}{2\tilde{\pi}} \right)^2 + \frac{1}{2\tilde{\pi}^2} \left(\sum_{\substack{k=-n \\ k \neq 0}}^n \frac{1}{k^2} - \sum_{\substack{k=-n \\ k \neq 0}}^n \frac{\cos(k(\tilde{\pi}+a))}{k^2} \right)$$


$$= \left(\frac{a + \tilde{\pi}}{2\tilde{\pi}} \right)^2 + \frac{1}{\tilde{\pi}^2} \left(\sum_{k=1}^n \frac{1}{k^2} - \sum_{k=1}^n \frac{\cos(k(\tilde{\pi}+a))}{k^2} \right)$$

General formula: $x \in [0, 2\tilde{\pi}]$

$$\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} = \frac{(x - \tilde{\pi})^2}{4} - \frac{\tilde{\pi}^2}{12}$$

$$\begin{array}{ccc}
 \downarrow n \rightarrow \infty & & \downarrow n \rightarrow \infty \\
 (*) & & (**) \\
 = \frac{\tilde{\pi}^2}{6} & & \frac{a^2}{4} - \frac{\tilde{\pi}^2}{12}
 \end{array}$$

$$\begin{aligned}
 \Rightarrow \sum_{k=-\infty}^{\infty} |c_k|^2 &= \left(\frac{a + \tilde{\pi}}{2\tilde{\pi}} \right)^2 + \frac{1}{\tilde{\pi}^2} \left(\frac{\tilde{\pi}^2}{6} - \frac{a^2}{4} + \frac{\tilde{\pi}^2}{12} \right) \\
 &= \left(\frac{a + \tilde{\pi}}{2\tilde{\pi}} \right)^2 + \frac{1}{4} - \frac{a^2}{4\tilde{\pi}^2} = \frac{2a\tilde{\pi} + \tilde{\pi}^2}{4\tilde{\pi}^2} + \frac{1}{4} \\
 &= \frac{a}{2\tilde{\pi}} + \frac{1}{2} = \frac{1}{2\tilde{\pi}} \cdot (a + \tilde{\pi}) = \frac{1}{2\tilde{\pi}} \int_{-\tilde{\pi}}^a 1 \, dx = \langle h_a, h_a \rangle \\
 &= \|h_a\|^2
 \end{aligned}$$



Fourier Transform - Part 11

Let's prove: $\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} = \frac{(x-\pi)^2}{4} - \frac{\pi^2}{12}, \quad x \in [0, 2\pi]$

Note: $\frac{1}{2} + \sum_{k=1}^n \cos(kx) = \frac{1}{2} + \sum_{k=1}^n \frac{1}{2} \cdot (e^{ikx} + e^{-ikx}) = \frac{1}{2} \sum_{k=-n}^n e^{ikx}$

$$= \frac{1}{2} e^{-inx} \sum_{k=0}^{2n} \underbrace{e^{ikx}}_{q^k} \quad \text{with } q = e^{ix}$$
$$= \frac{1}{2} e^{-inx} \cdot \frac{1 - q^{2n+1}}{1 - q} \quad \text{geometric sum formula } q \neq 1$$
$$= \frac{1}{2} \frac{e^{-inx} - e^{i(n+1)x}}{1 - e^{ix}} \cdot \frac{-e^{-\frac{1}{2}ix}}{-e^{-\frac{1}{2}ix}}$$
$$= \frac{1}{2} \frac{e^{i(n+\frac{1}{2})x} - e^{-i(n+\frac{1}{2})x}}{e^{\frac{1}{2}ix} - e^{-\frac{1}{2}ix}} \cdot \frac{\frac{1}{2i}}{\frac{1}{2i}} = \frac{1}{2} \frac{\sin((n+\frac{1}{2})x)}{\sin(\frac{1}{2}x)}$$

for $x \in \mathbb{R} \setminus \{2\pi m \mid m \in \mathbb{Z}\}$

Lemma: $\sum_{k=1}^{\infty} \frac{\sin(kx)}{k} = \frac{\pi - x}{2} \quad \text{for } x \in (0, 2\pi)$

and we have uniform convergence on interval $[\varepsilon, 2\pi - \varepsilon], \varepsilon > 0$.

Proof: $\sum_{k=1}^n \frac{\sin(kx)}{k} = \sum_{k=1}^n \int_{\pi}^x \cos(kt) dt = \int_{\pi}^x \sum_{k=1}^n \cos(kt) dt$

$$= \int_{\pi}^x \left(\frac{1}{2} \frac{\sin((n+\frac{1}{2})t)}{\sin(\frac{1}{2}t)} - \frac{1}{2} \right) dt$$
$$= \int_{\pi}^x \underbrace{\frac{\sin((n+\frac{1}{2})t)}{2 \sin(\frac{1}{2}t)}}_{f_n(x)} dt - \frac{1}{2}(x - \pi)$$

integration by parts: $f_n(x) = \int_{\pi}^x \underbrace{\frac{1}{2 \sin(\frac{1}{2}t)}}_u \cdot \underbrace{\sin((n+\frac{1}{2})t)}_{v'} dt$


$u' = -\frac{1}{2} \frac{\cos(\frac{1}{2}t)}{(\sin(\frac{1}{2}t))^2}$

$v = \frac{1}{n+\frac{1}{2}} \cdot (-1) \cdot \cos((n+\frac{1}{2})t)$

$$f_n(x) = \frac{1}{n+\frac{1}{2}} \cdot \frac{(-1) \cos((n+\frac{1}{2})t)}{2 \sin(\frac{1}{2}t)} \Big|_{\pi}^x - \int_{\pi}^x \frac{1}{n+\frac{1}{2}} \frac{(-1) \cdot \cos((n+\frac{1}{2})t) \cos(\frac{1}{2}t)}{(-4) \cdot (\sin(\frac{1}{2}t))^2} dt$$


$$= \frac{1}{n+\frac{1}{2}} \left(\underbrace{\frac{(-1) \cos((n+\frac{1}{2})x)}{2 \sin(\frac{1}{2}x)}}_{= a(x)} - \frac{1}{4} \int_{\pi}^x \underbrace{\frac{\cos((n+\frac{1}{2})t) \cos(\frac{1}{2}t)}{(\sin(\frac{1}{2}t))^2}}_{b(x)} dt \right)$$

For $\epsilon > 0$, choose $x \in [\epsilon, 2\pi - \epsilon]$:



$$\|f_n\|_{\infty} \leq \frac{1}{n+\frac{1}{2}} \left(\|a\|_{\infty} + \|b\|_{\infty} \right)$$

$$\leq \frac{1}{n+\frac{1}{2}} \left(\frac{1}{2\delta} + \frac{1}{4\delta^2} \cdot \pi \right) \xrightarrow{n \rightarrow \infty} 0$$



Recall $f_n(x) = \sum_{k=1}^n \frac{\sin(kx)}{k} + \frac{1}{2}(x - \pi)$

□

Theorem: $\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} = \frac{(x-\pi)^2}{4} - \frac{\pi^2}{12}, \quad x \in [0, 2\pi]$

uniform convergence on $[0, 2\pi]$

Proof: For $\varepsilon > 0$, $x, x_0 \in [\varepsilon, 2\pi - \varepsilon]$: (use Lemma)

$$\int_{x_0}^x \sum_{k=1}^{\infty} \frac{\sin(kt)}{k} dt = \int_{x_0}^x \frac{\pi - t}{2} dt = -\frac{(\pi - t)^2}{4} \Big|_{x_0}^x = -\frac{(x - \pi)^2}{4} + \underbrace{\frac{(x_0 - \pi)^2}{4}}_{C_0}$$

uniform convergence \Rightarrow

$$\sum_{k=1}^{\infty} \int_{x_0}^x \frac{\sin(kt)}{k} dt = \sum_{k=1}^{\infty} -\frac{\cos(kt)}{k^2} \Big|_{x_0}^x = -\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} + C_1$$

$$\Rightarrow \sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} = \frac{(x - \pi)^2}{4} + C \quad \leftarrow \text{calculate it!}$$

\rightarrow still uniform convergence on $[\varepsilon, 2\pi - \varepsilon]$

We know more:

(1) $\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2}$ uniformly convergent on $[0, 2\pi]$

by Weierstrass M-test since $\left| \frac{\cos(kx)}{k^2} \right| \leq \frac{1}{k^2}$

$$\Rightarrow [0, 2\pi] \ni x \mapsto \sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} \text{ continuous function}$$

(2) $[0, 2\pi] \ni x \mapsto \frac{(x - \pi)^2}{4} + C$ continuous function


(3) $\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} = \frac{(x - \pi)^2}{4} + C$ for all $x \in (0, 2\pi)$

$$\Rightarrow \sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} = \frac{(x - \pi)^2}{4} + C \text{ uniformly convergent on } [0, 2\pi]$$


Find C : $\int_0^{2\pi} \sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} dx = \int_0^{2\pi} \left(\frac{(x - \pi)^2}{4} + C \right) dx = \underbrace{\frac{(x - \pi)^3}{12} \Big|_0^{2\pi}}_{\frac{\pi^3}{6}} + 2\pi \cdot C$

// uniform convergence

$$\sum_{k=1}^{\infty} \int_0^{2\pi} \frac{\cos(kx)}{k^2} dx = 0 \quad \Rightarrow \quad C = -\frac{\pi^2}{12}$$




Fourier Transform - Part 12



Parseval's identity holds for h_a for every possible a . (part 10)

Step functions: consider the complex vector space:

$$\mathcal{S}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) := \left\{ g \in \mathcal{F}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \mid \begin{array}{l} \text{there are } m \in \mathbb{N}, a_i \in [-\pi, \pi], \\ \lambda_i \in \mathbb{C} \text{ such that:} \\ g = \sum_{i=1}^m \lambda_i \cdot h_{a_i} \end{array} \right\}$$



Do we have Parseval's identity here?

Consider step function $g \in \mathcal{S}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \rightsquigarrow$ $g = \sum_{i=1}^m \lambda_i \cdot h_{a_i}$

$$c_k = \langle e_k, g \rangle = \left\langle e_k, \sum_{i=1}^m \lambda_i \cdot h_{a_i} \right\rangle = \sum_{i=1}^m \lambda_i \langle e_k, h_{a_i} \rangle$$

$$|c_k|^2 = \overline{c_k} c_k = \overline{\sum_{j=1}^m \lambda_j \langle e_k, h_{a_j} \rangle} \cdot \sum_{i=1}^m \lambda_i \langle e_k, h_{a_i} \rangle$$

$$= \sum_{j=1}^m \sum_{i=1}^m \overline{\lambda_j} \lambda_i \langle h_{a_j}, e_k \rangle \langle e_k, h_{a_i} \rangle$$

$$\sum_{k=-n}^n |c_k|^2 = \sum_{i,j=1}^m \overline{\lambda_j} \lambda_i \left(\sum_{k=-n}^n \langle h_{a_j}, e_k \rangle \langle e_k, h_{a_i} \rangle \right)$$

(part 9) $\xrightarrow{h \rightarrow \infty}$


informal:
 $\left(\sum_{k=-\infty}^{\infty} |e_k\rangle \langle e_k| = \mathbb{1} \right)$
 we have Parseval's identity for h_{a_j} and h_{a_i}

$$\langle h_{a_j}, h_{a_i} \rangle$$

$$\Rightarrow \sum_{k=-\infty}^{\infty} |c_k|^2 = \sum_{i,j=1}^m \overline{\lambda_j} \lambda_i \langle h_{a_j}, h_{a_i} \rangle = \left\langle \sum_{j=1}^m \lambda_j \cdot h_{a_j}, \sum_{i=1}^m \lambda_i \cdot h_{a_i} \right\rangle$$

$$= \langle g, g \rangle = \|g\|^2$$

Result: Parseval's identity holds for $\int_{2\pi\text{-per}} (\mathbb{R}, \mathbb{C}) \subseteq L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$.



Fourier Transform - Part 13

Theorem: $L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ with inner product $\langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \overline{f(x)} \cdot g(x) dx$

and ONS $(\dots, e_{-2}, e_{-1}, e_0, e_1, e_2, \dots)$ given by $e_k: x \mapsto e^{ikx}$.

For $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ define: $\mathcal{F}_n(f) = \sum_{k=-n}^n e_k \underbrace{\langle e_k, f \rangle}_{c_k}$.

Then: $\|f - \mathcal{F}_n(f)\| \xrightarrow{n \rightarrow \infty} 0$ L^2 -norm


(equivalent to Parseval's identity: $\|f\|^2 = \sum_{k=-\infty}^{\infty} |\langle e_k, f \rangle|^2$)

Fact: Continuous functions are dense in $L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$, which means:

For $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ and $\varepsilon > 0$, there is a 2π -periodic continuous function

$g: \mathbb{R} \rightarrow \mathbb{C}$ with $\|f - g\| < \varepsilon$. L^2 -norm

Proposition: $C_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ is dense in $L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$.




Proof: Let $\varepsilon > 0$, $f: [-\pi, \pi] \rightarrow \mathbb{C}$ square integrable.

Then there is a continuous function $g: [-\pi, \pi] \rightarrow \mathbb{C}$ with $\|f - g\| < \varepsilon$.

domain compact


$\Rightarrow g$ is uniformly continuous: for given $\varepsilon > 0$ there $\delta > 0$:

$$|x - y| < \delta \Rightarrow |g(x) - g(y)| < \varepsilon$$



Decompose $[-\pi, \pi]$: I_1, I_2, \dots, I_N

$\text{length}(I_j) < \delta$



We get: $|g(x) - h(x)| = |g(x) - g(y)|$ for $y \in \overline{I_j}$

\uparrow
 $x \in I_j$

$< \epsilon$ because $|x-y| < \delta$

In total: $\|f - h\| \leq \|f - g\| + \|g - h\| < \epsilon + C \cdot \epsilon$ □

$= \left(\int_{-\tilde{\pi}}^{\tilde{\pi}} \underbrace{|g(x) - h(x)|^2}_{< \epsilon} dx \right)^{1/2}$

Theorem (see above): For $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$: $\|f - \mathcal{F}_n(f)\| \xrightarrow{n \rightarrow \infty} 0$

Proof: Let $\epsilon > 0$, $f \in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$. Choose $h \in \mathcal{S}_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$ with $\|f - h\| < \epsilon$.

Then: $\|f - \mathcal{F}_n(f)\| = \|f + h - h - \mathcal{F}_n(f) + \mathcal{F}_n(h) - \mathcal{F}_n(h)\|$


$\leq \| (f - h) - \mathcal{F}_n(f - h) \| + \| h - \mathcal{F}_n(h) \|$

$\leq \|f - h\| < \epsilon$ $\xrightarrow{h \rightarrow \infty} 0$ (part 12)


Pythagorean theorem:

$\|(f - h) - \mathcal{F}_n(f - h)\|^2 + \|\mathcal{F}_n(f - h)\|^2 = \|f - h\|^2$

$\Rightarrow \lim_{n \rightarrow \infty} \|f - \mathcal{F}_n(f)\| = 0$ □



Fourier Transform - Part 14





$$\|f - \mathcal{F}_n(f)\|_{L^2} \xrightarrow{n \rightarrow \infty} 0$$

not a pointwise convergence!

→ We can get uniform convergence for special functions

Example: continuous and piecewise C^1 -function




Supremum norm:

$$\|f\|_{\infty} := \sup_{x \in [-\pi, \pi]} |f(x)|$$

$$\Rightarrow \int_{-\pi}^{\pi} |f(x)|^2 dx \leq \int_{-\pi}^{\pi} \|f\|_{\infty}^2 dx = 2\pi \cdot \|f\|_{\infty}^2$$

$$\Rightarrow \|f\|_{L^2} \leq \|f\|_{\infty}$$

Theorem: $f: \mathbb{R} \rightarrow \mathbb{C}$ 2π -periodic continuous function.




Assume there are finitely many points (a_1, a_2, \dots, a_m)

inside the interval $[-\pi, \pi]$ such that:

$$f|_{[a_j, a_{j+1}]} \in C^1 \quad \text{for all } j \in \{0, 1, \dots, m\}$$


Then: $\|f - \mathcal{F}_n(f)\|_\infty \xrightarrow{n \rightarrow \infty} 0$

$$\left(\begin{aligned} \mathcal{F}_n(f) &= \sum_{k=-n}^n e_k \langle e_k, f \rangle \\ e_k: x &\mapsto e^{ikx} \\ \langle f, g \rangle &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \overline{f(x)} \cdot g(x) dx \end{aligned} \right)$$



Fourier Transform - Part 15

Theorem: $f: \mathbb{R} \rightarrow \mathbb{C}$ 2π -periodic continuous function and piecewise C^1 -function:




there are finitely many points (a_1, a_2, \dots, a_m)

inside the interval $[-\pi, \pi]$ such that: $f|_{[a_j, a_{j+1}]} \in C^1$ for all $j \in \{0, 1, \dots, m\}$, $a_0 := -\pi$, $a_{m+1} := \pi$

Then: $\mathcal{F}_n(f) \xrightarrow{h \rightarrow \infty} f$ uniformly.

Proof: Consider the derivative function: $\tilde{f}(x) := \begin{cases} 0 & , x \in \{a_0, a_1, \dots, a_{m+1}\} \\ f'(x) & , \text{else} \end{cases}$



piecewise continuous function $\in L^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$

Parseval's identity: $\|\tilde{f}\|^2 = \sum_{k=-\infty}^{\infty} |\langle e_k, \tilde{f} \rangle|^2 < \infty$

What about the Fourier coefficients of f ? ($k \neq 0$)

$$c_k := \langle e_k, f \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \underbrace{e^{-ikx}}_{u'} \underbrace{f(x)}_v dx \stackrel{\text{integration by parts}}{=} \frac{1}{2\pi} \left(u \cdot v \Big|_{-\pi}^{\pi} - \int_{-\pi}^{\pi} u v' dx \right)$$

$$= \frac{1}{2\pi} \left(0 + \frac{1}{ik} \int_{-\pi}^{\pi} e^{-ikx} \tilde{f}(x) dx \right) = \frac{1}{ik} \langle e_k, \tilde{f} \rangle$$

General inequality for real numbers: $x \cdot y \leq \frac{x^2 + y^2}{2}$

$$|c_k| = \frac{1}{k} |\langle e_k, \tilde{f} \rangle| \leq \frac{1}{2} \left(\frac{1}{k^2} + |\langle e_k, \tilde{f} \rangle|^2 \right)$$

$$\sum_{\substack{k=-\infty \\ k \neq 0}}^{\infty} |C_k| \leq \sum_{\substack{k=-\infty \\ k \neq 0}}^{\infty} \frac{1}{k^2} + \sum_{\substack{k=-\infty \\ k \neq 0}}^{\infty} |\langle e_k, \tilde{f} \rangle|^2 < \infty$$

$$\mathcal{F}_n(f)(x) = \sum_{k=-n}^n \underbrace{e^{ikx} \cdot C_k}_{f_k(x)} \quad \text{with } |f_k(x)| \leq M_k =: |C_k|, \quad \sum_{k=-\infty}^{\infty} M_k < \infty$$

Weierstrass
M-Test

$$\implies \sum_{k=-\infty}^{\infty} f_k \quad \text{uniformly convergent to a continuous function}$$

$$h: [-\pi, \pi] \longrightarrow \mathbb{C}$$

Status quo: $\|\mathcal{F}_n(f) - h\|_{\infty} \xrightarrow{h \rightarrow \infty} 0, \quad \|\mathcal{F}_n(f) - f\|_{L^2} \xrightarrow{h \rightarrow \infty} 0$


More estimates: $\|f - h\|_{L^2} \leq \|f - \mathcal{F}_n(f)\|_{L^2} + \underbrace{\|\mathcal{F}_n(f) - h\|_{L^2}}_{\leq \|\mathcal{F}_n(f) - h\|_{\infty}}$

$$\xrightarrow{h \rightarrow \infty} 0$$


Hence: $\|f - h\|_{L^2} = 0 \xRightarrow{\text{continuous functions}} f = h$

Conclusion: $\|\mathcal{F}_n(f) - f\|_{\infty} \xrightarrow{h \rightarrow \infty} 0$ (uniform convergence of the Fourier series)

□



Fourier Transform - Part 16



⇒ continuous + piecewise C^1 -function

Example: $f: \mathbb{R} \rightarrow \mathbb{C}$ 2π -periodic with $f(x) = x^2 - \pi^2$ for $x \in [-\pi, \pi]$.

Let's calculate the Fourier coefficients: $c_k := \langle e_k, f \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikx} f(x) dx$

$$\begin{aligned} c_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} (x^2 - \pi^2) dx = \frac{1}{2\pi} \left(\frac{1}{3} x^3 - \pi^2 x \right) \Big|_{-\pi}^{\pi} \\ &= \frac{1}{2\pi} \cdot 2 \cdot \left(\frac{1}{3} \pi^3 - \pi^3 \right) = \underline{-\frac{2}{3} \pi^2} \end{aligned}$$

For $k \neq 0$: $c_k = \frac{1}{ik} \langle e_k, f' \rangle$ (integration by parts, see part 15)

$$\begin{aligned} &= \frac{1}{2\pi ik} \int_{-\pi}^{\pi} \underbrace{e^{-ikx}}_u \cdot \underbrace{2x}_v dx \quad (\text{integration by parts}) \\ &\quad \rightarrow u = \frac{1}{-ik} e^{-ikx}, \quad v' = 2 \\ &= \frac{1}{2\pi ik} \left(-\frac{1}{ik} e^{-ikx} \cdot 2x \Big|_{-\pi}^{\pi} - \int_{-\pi}^{\pi} \underbrace{\left(-\frac{1}{ik} e^{-ikx} \right) \cdot 2}_{=0} dx \right) \\ &= \frac{1}{\pi \cdot k^2} \left(\underbrace{e^{-ik\pi}}_{=(-1)^k} \pi - \underbrace{e^{ik\pi}}_{=(-1)^k} (-\pi) \right) \\ &= \frac{2 \cdot (-1)^k}{k^2} \end{aligned}$$

Fourier series: $x^2 - \pi^2 = \sum_{k=-\infty}^{\infty} C_k e^{ikx} = -\frac{2}{3}\pi^2 + \sum_{\substack{k=-\infty \\ k \neq 0}}^{\infty} \frac{2 \cdot (-1)^k}{k^2} \underbrace{e^{ikx}}_{\cos(kx) + i \cdot \sin(kx)}$

$x \in [-\pi, \pi]$

$$= -\frac{2}{3}\pi^2 + 2 \cdot \sum_{k=1}^{\infty} \frac{2 \cdot (-1)^k}{k^2} \cos(kx)$$

For all $x \in [-\pi, \pi]$: $x^2 - \frac{1}{3}\pi^2 = \sum_{k=1}^{\infty} \frac{4}{k^2} (-1)^k \cos(kx)$ ← uniform convergence!


In particular for $x=0$: $-\frac{1}{3}\pi^2 = \sum_{k=1}^{\infty} \frac{4}{k^2} (-1)^k$

$$\Rightarrow \sum_{k=1}^{\infty} \frac{(-1)^k}{k^2} = -\frac{1}{12}\pi^2$$

Parseval's identity: $\sum_{k=-\infty}^{\infty} |C_k|^2 = \|f\|_{L^2}^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} (x^2 - \pi^2)^2 dx = \frac{8}{15}\pi^4$


$$|C_0|^2 + \sum_{\substack{k=-\infty \\ k \neq 0}}^{\infty} \left| \frac{2 \cdot (-1)^k}{k^2} \right|^2$$

$$\frac{4}{9} \cdot \pi^4 + 2 \cdot \sum_{k=1}^{\infty} \frac{4}{k^4} \Rightarrow \sum_{k=1}^{\infty} \frac{1}{k^4} = \frac{\pi^4}{90}$$



Fourier Transform - Part 17


$$\begin{array}{l}
 f: \mathbb{R} \rightarrow \mathbb{C} \quad 2\pi\text{-periodic} \\
 f \in \mathcal{L}^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C}) \\
 \text{continuous + piecewise } C^1\text{-function}
 \end{array}
 \implies
 \begin{array}{l}
 \mathcal{F}_n(f) \xrightarrow{h \rightarrow \infty} f \quad (\text{in } L^2\text{-norm}) \\
 \mathcal{F}_n(f) \xrightarrow{h \rightarrow \infty} f \quad (\text{pointwisely}) \\
 \mathcal{F}_n(f) \xrightarrow{h \rightarrow \infty} f \quad (\text{uniformly})
 \end{array}$$




Theorem: $f \in \mathcal{L}^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$, $\hat{x} \in [-\pi, \pi]$ with:

$$\begin{array}{ll}
 f(\hat{x}^-) := \lim_{\substack{\epsilon \rightarrow 0 \\ \epsilon > 0}} f(\hat{x} - \epsilon) \text{ exists,} & \lim_{\substack{h \rightarrow 0 \\ h < 0}} \frac{f(\hat{x} + h) - f(\hat{x})}{h} \text{ exists} \\
 f(\hat{x}^+) := \lim_{\substack{\epsilon \rightarrow 0 \\ \epsilon > 0}} f(\hat{x} + \epsilon) \text{ exists,} & \lim_{\substack{h \rightarrow 0 \\ h > 0}} \frac{f(\hat{x} + h) - f(\hat{x})}{h} \text{ exists}
 \end{array}$$

Then: $\mathcal{F}_n(f)(\hat{x}) \xrightarrow{h \rightarrow \infty} \frac{1}{2} \left(f(\hat{x}^+) + f(\hat{x}^-) \right)$



Example:



$$f(x) = \begin{cases} 0 & , x \in [-\pi, 0) \\ \pi - x & , x \in [0, \pi) \end{cases}$$

Fourier coefficients: $C_k := \langle e_k, f \rangle = \frac{1}{2\pi} \int_0^{\pi} e^{-ikx} (\pi - x) dx$


$$= \begin{cases} \frac{\pi}{4} & , k = 0 \\ \frac{1}{2\pi} \cdot \left(-\frac{1}{k^2}((-1)^k - 1) - i \frac{\pi}{k} \right) & , k \neq 0 \end{cases}$$

Fourier series:

$$\mathcal{F}_n(f)(x) = \frac{\pi}{4} + \sum_{\substack{k=-n \\ k \neq 0}}^n C_k \cdot e^{ikx}$$

$$a_k = C_k + C_{-k}$$

$$b_k = i \cdot (C_k - C_{-k})$$




$$= \frac{\pi}{4} + \sum_{k=1}^n a_k \cos(kx) + \sum_{k=1}^n b_k \sin(kx)$$

$\underbrace{a_k}_{\frac{1}{\pi} \cdot \frac{1}{k^2} (1 - (-1)^k)} \quad \underbrace{b_k}_{\frac{1}{k}}$

$$\mathcal{F}_n(f)(0) \xrightarrow{n \rightarrow \infty} \frac{\pi}{2}$$

$$\Rightarrow \frac{\pi}{2} = \frac{\pi}{4} + \sum_{k=1}^{\infty} \frac{1}{\pi} \frac{1 - (-1)^k}{k^2}$$

$$\Rightarrow \frac{\pi^2}{4} = \sum_{k=1}^{\infty} \frac{1 - (-1)^k}{k^2}$$




Fourier Transform - Part 18

Definition: The continuous function $\mathcal{D}_n: \mathbb{R} \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, given by

$$\mathcal{D}_n(x) = \sum_{k=-n}^n e^{ikx} \stackrel{\text{part 11}}{=} 1 + 2 \cdot \sum_{k=1}^n \cos(kx) \stackrel{\text{part 11}}{=} \frac{\sin\left(\left(n+\frac{1}{2}\right)x\right)}{\sin\left(\frac{1}{2}x\right)}$$

is called the Dirichlet kernel.

$2\pi \cdot m$ zeros
for $m \in \mathbb{Z}$




2π -periodic

For Fourier series:

$$\begin{aligned} \mathcal{F}_n(f)(x) &= \sum_{k=-n}^n c_k \cdot e^{ikx} = \sum_{k=-n}^n \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-iky} f(y) dy \right) \cdot e^{ikx} \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) \sum_{k=-n}^n e^{ik(x-y)} dy = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) \mathcal{D}_n(\underbrace{x-y}_z) dy \\ &= \frac{1}{2\pi} \int_{x-\pi}^{x+\pi} f(x-z) \mathcal{D}_n(z) dz = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathcal{D}_n(z) f(x-z) dz \\ &= \langle \mathcal{D}_n, f(x-\cdot) \rangle = \frac{1}{2\pi} (\mathcal{D}_n * f)(x) \quad (\text{convolution}) \end{aligned}$$

Properties: (1) \mathcal{D}_n has exactly $2n$ zeros inside the interval $[-\pi, \pi]$




$$\frac{\sin\left(\left(n+\frac{1}{2}\right)x\right)}{\sin\left(\frac{1}{2}x\right)}$$


$$(2) \int_{-\pi}^{\pi} \mathcal{D}_n(x) dx = \int_{-\pi}^{\pi} (1 + e^{ix} + e^{-ix} + e^{2ix} + e^{-2ix} + \dots + e^{nix} + e^{-nix}) dx$$

$$= 2\pi \quad \Rightarrow \quad \langle \mathcal{D}_n, 1 \rangle = 1$$

$$(3) \int_{-\pi}^{\pi} |\mathcal{D}_n(x)| dx \xrightarrow{h \rightarrow \infty} \infty$$



Proof of (3): $|\mathcal{D}_n(x)| = \frac{|\sin((n+\frac{1}{2})x)|}{|\sin(\frac{1}{2}x)|}$



$$\geq \frac{|\sin((n+\frac{1}{2})x)|}{x} \quad \text{for all } x > 0$$

$$\int_{-\pi}^{\pi} |\mathcal{D}_n(x)| dx = 2 \cdot \int_0^{\pi} |\mathcal{D}_n(x)| dx \geq 2 \cdot \int_0^{\pi} \frac{|\sin((n+\frac{1}{2})x)|}{x} dx$$

$$= 2 \cdot \int_0^{(n+\frac{1}{2})\pi} \frac{|\sin(y)|}{y} dy \geq 2 \cdot \int_0^{n\pi} \frac{|\sin(y)|}{y} dy$$

$$= 2 \cdot \sum_{k=1}^h \int_{(k-1)\pi}^{k\pi} \frac{|\sin(y)|}{y} dy$$


maximal $k \cdot \pi$

$$\geq 2 \cdot \sum_{k=1}^h \int_{(k-1)\pi}^{k\pi} \frac{|\sin(y)|}{k\pi} dy$$

$$= 2 \cdot \sum_{k=1}^h \frac{1}{k\pi} \underbrace{\int_{(k-1)\pi}^{k\pi} |\sin(y)| dy}_{=1} = \text{const} \cdot \sum_{k=1}^h \frac{1}{k}$$


$$\xrightarrow{h \rightarrow \infty} \infty$$

$$\frac{|\sin(y)|}{y}$$



Fourier Transform - Part 19

Theorem: $f \in \mathcal{L}^2_{2\pi\text{-per}}(\mathbb{R}, \mathbb{C})$, $\hat{x} \in [-\pi, \pi]$ with:




$$f(\hat{x}^-) := \lim_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} f(\hat{x} - \varepsilon) \text{ exists, } \lim_{\substack{h \rightarrow 0 \\ h < 0}} \frac{f(\hat{x} + h) - f(\hat{x})}{h} \text{ exists}$$

$$f(\hat{x}^+) := \lim_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} f(\hat{x} + \varepsilon) \text{ exists, } \lim_{\substack{h \rightarrow 0 \\ h > 0}} \frac{f(\hat{x} + h) - f(\hat{x})}{h} \text{ exists}$$

Then: $\mathcal{F}_n(f)(\hat{x}) \xrightarrow{n \rightarrow \infty} \frac{1}{2} (f(\hat{x}^+) + f(\hat{x}^-)) =: M$

Proof: Dirichlet kernel: $\mathcal{D}_n(x) = \frac{\sin((n+\frac{1}{2})x)}{\sin(\frac{1}{2}x)}$ gives $\mathcal{F}_n(f)(\hat{x}) = \langle \mathcal{D}_n, f(\hat{x} - \cdot) \rangle$

and $\langle \mathcal{D}_n, M \rangle = M$



Use symmetry: $\langle \mathcal{D}_n, f(\hat{x} - \cdot) \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathcal{D}_n(x) f(\hat{x} - x) dx$

$$\begin{aligned} &= \frac{1}{2\pi} \left(\int_{-\pi}^0 \mathcal{D}_n(x) f(\hat{x} - \underbrace{x}_y) dx + \int_0^{\pi} \mathcal{D}_n(x) f(\hat{x} - x) dx \right) \\ &= \frac{1}{2\pi} \left(\int_0^{\pi} \mathcal{D}_n(y) f(\hat{x} + y) dy + \int_0^{\pi} \mathcal{D}_n(x) f(\hat{x} - x) dx \right) \\ &= \frac{1}{2\pi} \int_0^{\pi} \mathcal{D}_n(y) (f(\hat{x} + y) + f(\hat{x} - y)) dy \end{aligned}$$

Pointwise limit: $\mathcal{F}_n(f)(\hat{x}) - M = \langle \mathcal{D}_n, f(\hat{x}-\cdot) \rangle - \langle \mathcal{D}_n, M \rangle$

$$= \frac{1}{2\pi} \int_0^{\pi} \mathcal{D}_n(y) (f(\hat{x}+y) + f(\hat{x}-y)) dy - \frac{1}{2\pi} \int_0^{\pi} \mathcal{D}_n(y) \underbrace{2 \cdot M}_{f(\hat{x}^+) + f(\hat{x}^-)} dy$$

$$= \frac{1}{2\pi} \int_0^{\pi} \mathcal{D}_n(y) (f(\hat{x}+y) - f(\hat{x}^+) + f(\hat{x}-y) - f(\hat{x}^-)) dy$$

$$= \frac{1}{2\pi} \int_0^{\pi} \sin((n+\frac{1}{2})y) \underbrace{\frac{f(\hat{x}+y) - f(\hat{x}^+) + f(\hat{x}-y) - f(\hat{x}^-)}{\sin(\frac{1}{2}y)}}_{g(y)} dy$$

In the case that $g \in L^2_{2\pi\text{-per}}$, we get:

$$\frac{1}{2\pi} \int_0^{\pi} \sin((n+\frac{1}{2})y) g(y) dy$$

$$\left(\frac{1}{2i} (e^{iny} e^{i\frac{1}{2}y} - e^{-iny} e^{-i\frac{1}{2}y}) \right)$$

$$\langle e_{-n}, g_1 \rangle + \langle e_n, g_2 \rangle \xrightarrow[n \rightarrow \infty]{\text{part 8}} 0$$


(Bessel's inequality)

$L^2_{2\pi\text{-per}}$ -functions

Show that $g \in L^2_{2\pi\text{-per}}$:

$$g(y) = \begin{cases} \frac{f(\hat{x}+y) - f(\hat{x}^+)}{\sin(\frac{1}{2}y)} + \frac{f(\hat{x}-y) - f(\hat{x}^-)}{\sin(\frac{1}{2}y)}, & y \in (0, \pi) \\ 0, & y \in [-\pi, 0] \end{cases}$$

Does $g(y)$ explode for $y \rightarrow 0^+$?



$$\Rightarrow \left| \frac{f(\hat{x}+y) - f(\hat{x}^+)}{\sin(\frac{1}{2}y)} \right| \leq 4 \cdot \left| \frac{f(\hat{x}+y) - f(\hat{x}^+)}{y} \right|$$

$$\xrightarrow{y \rightarrow 0^+} 4 \cdot |C^+|$$

because $\lim_{\substack{h \rightarrow 0 \\ h > 0}} \frac{f(\hat{x}+h) - f(\hat{x})}{h} =: C^+$

and $\left| \frac{f(\hat{x}-y) - f(\hat{x}^-)}{\sin(\frac{1}{2}y)} \right| \leq 4 \cdot \left| \frac{f(\hat{x}-y) - f(\hat{x}^-)}{y} \right| \xrightarrow{y \rightarrow 0^+} 4 \cdot |C^-|$

because $\lim_{\substack{h \rightarrow 0 \\ h < 0}} \frac{f(\hat{x}+h) - f(\hat{x})}{h} =: C^-$

□