


Linear Algebra - Part 5



$$\mathbb{R}^n = \underbrace{\mathbb{R} \times \cdots \times \mathbb{R}}_{n \text{ times}} \quad \text{for } n \in \mathbb{N}$$

write $v \in \mathbb{R}^n$ in column form: $v = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} \in \mathbb{R}^n$

addition: $u + v = \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix} + \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} := \begin{pmatrix} u_1 + v_1 \\ \vdots \\ u_n + v_n \end{pmatrix}$

scalar multiplication: $\lambda \cdot u = \lambda \cdot \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix} := \begin{pmatrix} \lambda \cdot u_1 \\ \vdots \\ \lambda \cdot u_n \end{pmatrix}$

$\hookrightarrow (\mathbb{R}^n, +, \cdot)$ is a vector space

Properties: (a) $(\mathbb{R}^n, +)$ is an abelian group:

$$(1) \quad u + (v + w) = (u + v) + w \quad (\text{associativity of } +)$$

$$(2) \quad v + 0 = v \quad \text{with} \quad 0 = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix} \quad (\text{neutral element})$$

$$(3) \quad v + (-v) = 0 \quad \text{with} \quad -v = \begin{pmatrix} -v_1 \\ \vdots \\ -v_n \end{pmatrix} \quad (\text{inverse elements})$$

$$(4) \quad v + w = w + v \quad (\text{commutativity of } +)$$

(b) scalar multiplication is compatible: $\cdot : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$

$$(5) \quad \lambda \cdot (\mu \cdot v) = (\lambda \cdot \mu) \cdot v$$

$$(6) \quad 1 \cdot v = v$$

(c) distributive laws:

$$(7) \quad \lambda \cdot (v + w) = \lambda \cdot v + \lambda \cdot w$$

$$(8) \quad (\lambda + \mu) \cdot v = \lambda \cdot v + \mu \cdot v$$

Canonical unit vectors:

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \quad \dots, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

$v = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} \in \mathbb{R}^n$ can be written as a linear combination: $v = \sum_{j=1}^n v_j \cdot e_j$