## Linear Algebra - Part 12

Example: Xavier is two years older than Yasmin.

Together they are 40 years old.

How old is Xavier? How old is Yasmin?


Another Example: 
$$2 \times_{1} - 3 \times_{2} + 4 \times_{3} = -7$$
  
 $-3 \times_{1} + \times_{2} - \times_{3} = 0$   
 $20 \times_{1} + 10 \times_{2} = 80$   
 $10 \times_{2} + 25 \times_{3} = 90$ 
4 equations and 3 unknowns  $X_{1}, X_{2}, X_{3}$ 

Linear equation: constant  $X_1$  + constant  $X_2$  + ... + constant  $X_n$  = constant


Definition: System of linear equations (LES) with m equations and n unknowns:

A solution of the LES: choice of values for  $X_1, ..., X_n$  such that all mequations are satisfied.


Note: - it's possible that there is no solution m = 2, n = 2



- it's possible that there is a unique solution m = 2, n = 2



- it's possible that there are infinitely many solutions



Short notation: Instead of

$$a_{11} X_1 + a_{12} X_2 + \cdots + a_{1n} X_n = b_1$$

$$\alpha_{21} \times_1 + \alpha_{22} \times_2 + \cdots + \alpha_{2n} \times_n = b_2$$

$$a_{m1} \times_1 + a_{m2} \times_2 + \cdots + a_{mn} \times_n = b_m$$

we write

$$A \times = b$$

with 
$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$
,  $b = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}$ 

and 
$$X = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}$$

Example:

$$2 x_{1} - 3 x_{1} + 4 x_{3} = -7$$

$$-3 x_{1} + x_{2} - x_{3} = 0$$

$$20 x_{1} + 10 x_{2} = 80$$

$$10 x_{2} + 25 x_{3} = 90$$
can be written as

$$\begin{pmatrix} 2 & -3 & 4 \\ -3 & 1 & -1 \\ 20 & 10 & 0 \\ 0 & 10 & 25 \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -7 \\ 0 \\ 80 \\ 90 \end{pmatrix}$$

matrix-vector product

"matrix times vector = vector"