



## Linear Algebra - Part 26



dimension = 2



dimension = 1

### Steinitz Exchange Lemma

Let  $U \subseteq \mathbb{R}^n$  be a subspace and

$\mathcal{B} = (v^{(1)}, v^{(2)}, \dots, v^{(k)})$  be a basis of  $U$ .

$\mathcal{A} = (a^{(1)}, a^{(2)}, \dots, a^{(l)})$  linearly independent vectors in  $U$ .



Then: One can add  $k-l$  vectors from  $\mathcal{B}$  to the family  $\mathcal{A}$  such that we get a new basis of  $U$ .

Proof:  $l=1$  :  $\mathcal{B} \cup \mathcal{A} = (v^{(1)}, v^{(2)}, \dots, v^{(k)}, a^{(1)})$  is linearly dependent

because  $\mathcal{B}$  is a basis: there are uniquely given  $\lambda_1, \dots, \lambda_k \in \mathbb{R}$ :

$$(*) \quad a^{(1)} = \lambda_1 v^{(1)} + \dots + \lambda_k v^{(k)} \quad \text{Diagram: a green vector } a^{(1)} \text{ is shown as a linear combination of red vectors } v^{(1)}, \dots, v^{(k)}.$$

Choose  $\lambda_j \neq 0$  :

$$v^{(j)} = \frac{1}{\lambda_j} \left( \lambda_1 v^{(1)} + \dots + \lambda_{j-1} v^{(j-1)} + \lambda_{j+1} v^{(j+1)} + \dots + \lambda_k v^{(k)} - a^{(1)} \right)$$

Remove  $v^{(j)}$  from  $\mathcal{B} \cup \mathcal{A}$  and call it  $\mathcal{C}$ .

$\mathcal{E}$  is linearly independent:

$$\tilde{\lambda}_1 v^{(1)} + \dots + \tilde{\lambda}_{j-1} v^{(j-1)} + \tilde{\lambda}_j a^{(1)} + \tilde{\lambda}_{j+1} v^{(j+1)} + \dots + \tilde{\lambda}_k v^{(k)} = 0$$

Assume  $\tilde{\lambda}_j \neq 0$ :  $a^{(1)}$  = linear combination with  $v^{(1)}, \dots, v^{(j-1)}, v^{(j+1)}, \dots, v^{(k)}$

Hence:  $\tilde{\lambda}_j = 0 \Rightarrow$   $\Downarrow (*)$

$$\tilde{\lambda}_1 v^{(1)} + \dots + \tilde{\lambda}_{j-1} v^{(j-1)} + \tilde{\lambda}_{j+1} v^{(j+1)} + \dots + \tilde{\lambda}_k v^{(k)} = 0$$

lin. independence

$$\Rightarrow \tilde{\lambda}_i = 0 \text{ for } i \in \{1, \dots, k\}$$

$\mathcal{E}$  spans  $U$ :  $u \in U \stackrel{\mathcal{B} \text{ basis}}{\Rightarrow}$  there are coefficients

$$v^{(j)} = \frac{1}{\tilde{\lambda}_j} (\tilde{\lambda}_1 v^{(1)} + \dots + \tilde{\lambda}_{j-1} v^{(j-1)} + \tilde{\lambda}_{j+1} v^{(j+1)} + \dots + \tilde{\lambda}_k v^{(k)} - a^{(1)})$$

$$u = \mu_1 v^{(1)} + \dots + \mu_{j-1} v^{(j-1)} + \mu_j v^{(j)} + \mu_{j+1} v^{(j+1)} + \dots + \mu_k v^{(k)}$$

$$= \tilde{\mu}_1 v^{(1)} + \dots + \tilde{\mu}_{j-1} v^{(j-1)} + \tilde{\mu}_j a^{(1)} + \tilde{\mu}_{j+1} v^{(j+1)} + \dots + \tilde{\mu}_k v^{(k)}$$