



## Linear Algebra - Part 41

$A \in \mathbb{R}^{m \times h}$   $\xrightarrow{\text{Gaussian elimination}}$  row echelon form

$$\left( \begin{array}{ccccc|c} x_1 & x_2 & x_3 & x_4 & x_5 & \\ \hline 1 & 2 & 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & -1 & 4 & 0 \\ 0 & 0 & 0 & 4 & 8 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

$$\Rightarrow \text{Ker}(A) = \left\{ x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + x_5 \begin{pmatrix} 2 \\ 0 \\ -3 \\ -2 \\ 1 \end{pmatrix} \mid x_2, x_5 \in \mathbb{R} \right\}$$

Remember:

$$\begin{aligned} \dim(\text{Ker}(A)) &= \text{number of free variables} \\ + \\ \dim(\text{Ran}(A)) &= \text{number of leading variables} \\ &= h \end{aligned}$$

Proposition: For  $A \in \mathbb{R}^{m \times h}$  and  $b \in \mathbb{R}^m$ , we have the following equivalences:

- (1)  $Ax = b$  has at least one solution.
- (2)  $b \in \text{Ran}(A)$
- (3)  $b$  can be written as a linear combination of the columns of  $A$ .

(4) Row echelon form looks like:

$$\left( \begin{array}{cccc|c} \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \vdots & \dots & \dots & \dots & \vdots \\ 0 & \dots & \dots & \dots & 0 \end{array} \right)$$

Proof: (1)  $\Leftrightarrow$  (2) given by definition of  $\text{Ran}(A)$

(2)  $\Leftrightarrow$  (3) given by column picture of  $\text{Ran}(A)$

$$\begin{aligned}\text{Ran}(A) &= \left\{ \left( \begin{array}{c|c} | & | \\ a_1 & \dots & a_n \\ | & & | \end{array} \right) x \mid x \in \mathbb{R}^n \right\} \\ &= \left\{ x_1 \begin{pmatrix} | \\ a_1 \\ | \end{pmatrix} + \dots + x_n \begin{pmatrix} | \\ a_n \\ | \end{pmatrix} \mid x \in \mathbb{R}^n \right\}\end{aligned}$$

(4)  $\Rightarrow$  (1)

Assume we have this:  $\left( \begin{array}{cccc|c} \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ 0 & \dots & 0 & 0 & 0 \\ \vdots & & \vdots & \vdots & \vdots \\ 0 & \dots & 0 & 0 & 0 \end{array} \right)$

Then solve  $\left( \begin{array}{cccc|c} \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ 0 & \dots & 0 & 0 & 0 \\ \vdots & & \vdots & \vdots & \vdots \\ 0 & \dots & 0 & 0 & 0 \end{array} \right)$  by backwards substitution.

(or argue with  $\text{rank}(A) = \text{rank}((A|b))$ )

(1)  $\Rightarrow$  (4) (let's show:  $\neg(4) \Rightarrow \neg(1)$ )

Assume:  $\left( \begin{array}{cccc|c} \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} & \text{---} \\ 0 & \dots & 0 & 0 & c \\ \vdots & & \vdots & \vdots & \vdots \\ 0 & \dots & 0 & 0 & c \end{array} \right)$   $\rightarrow$  not solvable  $0 = c \neq 0$   
 $\Rightarrow$  no solution for  $Ax = b$   $\square$