

Linear Algebra - Part 52

We know for
$$A \in \mathbb{R}^{2 \times 2}$$
: $\det(A) \neq 0 \iff A_{X} = b$ has a unique solution

For
$$A \in \mathbb{R}^{n \times n}$$
: $det(A) = 0 \iff A$ singular

Proposition: For $A \in \mathbb{R}^{n \times n}$, the following claims are equivalent:

- $det(A) \neq 0$
- · columns of A are linearly independent
- · rows of A are linearly independent
- rank(A) = h
- Ker(A) = {0}
- · A is invertible
- Ax = b has a unique solution for each $b \in \mathbb{R}^n$

Cramer's rule: $A \in \mathbb{R}^{n \times n}$ non-singular, $b \in \mathbb{R}^n$, $x = \begin{pmatrix} x_1 \\ x_2 \\ x_n \end{pmatrix} \in \mathbb{R}^n$ unique solution of Ax = b.

Then:
$$X_{i} = \frac{\det \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0_{1} & \cdots & 0_{i-1} & b & a_{i+1} & \cdots & a_{h} \\ 1 & \cdots & a_{i-1} & a_{i} & a_{i+1} & \cdots & a_{h} \\ 1 & \cdots & a_{i-1} & a_{i} & a_{i+1} & \cdots & a_{h} \end{pmatrix}$$

Laplace expansion
$$= det \begin{pmatrix} a_1 \cdots a_{j-1} & e_i & a_{j+1} \cdots a_h \end{pmatrix}$$

We can show:
$$A^{-1} = \frac{C_1^T}{\det(A)}$$

Hence:
$$X = A^{-1}L = \frac{C_1^{-1}L}{\det(A)}$$
 and

$$X = \overline{A^1 l} = \frac{C_1^T l}{\det(A)} \quad \text{and} \quad (C_1^T l)_i = \sum_{k=1}^n (C_1^T)_{ik} l_k = \sum_{k=1}^n C_{ki} l_k$$

linear in the ith column
$$= \det \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix}$$