


## Linear Algebra - Part 53

eigenvalue (German: Eigenwert) (David Hilbert, 1904)

↳ proper/own/characteristic

Consider:  $A \in \mathbb{R}^{n \times n} \iff f_A: \mathbb{R}^n \rightarrow \mathbb{R}^n$  linear map



Question: Are there vectors which are only scaled by  $f_A$ ?

Answer:  $Ax = \lambda \cdot x$  for a number  $\lambda \in \mathbb{R}$

$$\iff (A - \lambda \mathbb{1})x = 0 \quad \text{for a number } \lambda \in \mathbb{R}$$

$$\iff x \in \text{Ker}(A - \lambda \mathbb{1}) \quad \text{for a number } \lambda \in \mathbb{R}$$

↖ eigenvalue (if  $x \neq 0$ )      ↗ eigenvalue

Example:

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \lambda \cdot \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \iff \begin{array}{l} x_1 + x_2 = \lambda \cdot x_1 \quad \text{I} \\ x_2 = \lambda \cdot x_2 \quad \text{II} \end{array}$$

For II:  $\lambda = 1$  or  $x_2 = 0$

$$\implies x_1 = \lambda \cdot x_1 \implies \lambda = 1 \text{ or } x_1 = 0$$

For I:  $x_1 + x_2 = x_1 \implies x_2 = 0$

solution: eigenvalue:  $\lambda = 1$

eigenvectors:  $x = \begin{pmatrix} x_1 \\ 0 \end{pmatrix}$  for  $x_1 \in \mathbb{R} \setminus \{0\}$

Definition:  $A \in \mathbb{R}^{n \times n}$ ,  $\lambda \in \mathbb{R}$ .

If there is  $x \in \mathbb{R}^n \setminus \{0\}$  with  $Ax = \lambda x$ , then:

- $\lambda$  is called an eigenvalue of  $A$
- $x$  is called an eigenvector of  $A$  (associated to  $\lambda$ )
- $\text{Ker}(A - \lambda \mathbb{1})$  eigenspace of  $A$  (associated to  $\lambda$ )

The set of all eigenvalues of  $A$ :  $\text{spec}(A)$  spectrum of  $A$