


Linear Algebra - Part 54


$$A \in \mathbb{R}^{n \times n} \iff f_A: \mathbb{R}^n \longrightarrow \mathbb{R}^n \quad \text{linear map}$$

$$\text{eigenvalue equation: } Ax = \lambda \cdot x, \quad x \neq 0$$

optimal coordinate system: $A \in \mathbb{R}^{2 \times 2}, \quad Ax = 2x, \quad Ay = 1y$



$$u = a \cdot x + b \cdot y$$



$$\begin{aligned} Au &= A(a \cdot x + b \cdot y) \\ &= a \cdot Ax + b \cdot Ay \\ &= 2ax + 1by \end{aligned}$$

How to find enough eigenvectors?

$$x \neq 0 \text{ eigenvector associated to eigenvalue } \lambda \iff x \in \text{Ker}(A - \lambda \mathbb{1})$$

singular matrix

$$\det(A - \lambda \mathbb{1}) = 0 \iff \text{Ker}(A - \lambda \mathbb{1}) \text{ is non-trivial}$$

$$\iff \lambda \text{ is eigenvalue of } A$$

Example:

$$A = \begin{pmatrix} 3 & 2 \\ 1 & 4 \end{pmatrix}, \quad A - \lambda \mathbb{1} = \begin{pmatrix} 3-\lambda & 2 \\ 1 & 4-\lambda \end{pmatrix}$$

$$\det \begin{pmatrix} 3-\lambda & 2 \\ 1 & 4-\lambda \end{pmatrix} = (3-\lambda)(4-\lambda) - 2 \quad \text{characteristic polynomial}$$

$$= 10 - 7\lambda + \lambda^2$$

$$= (\lambda - 5)(\lambda - 2) \stackrel{!}{=} 0$$

$$\Rightarrow 2 \text{ and } 5 \text{ are eigenvalues of } A$$

General case: For $A \in \mathbb{R}^{n \times n}$:

$$\det(A - \lambda \mathbb{1}) = \det \begin{pmatrix} a_{11} - \lambda & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} - \lambda & & \vdots \\ \vdots & & \ddots & \\ a_{n1} & \cdots & & a_{nn} - \lambda \end{pmatrix}$$

Leibniz formula

$$\downarrow \\ = (a_{11} - \lambda) \cdots (a_{nn} - \lambda) + \cdots$$

$$= (-1)^n \cdot \lambda^n + c_{n-1} \lambda^{n-1} + \cdots + c_1 \lambda^1 + c_0$$

Definition: For $A \in \mathbb{R}^{n \times n}$, the polynomial of degree n given by

$$p_A: \lambda \mapsto \det(A - \lambda \mathbb{1})$$

is called the characteristic polynomial of A .

Remember: The zeros of the characteristic polynomial are exactly the eigenvalues of A .