


Linear Algebra - Part 64

Diagonalization = transform matrix into a diagonal one
= find a an optimal coordinate system


Example:

$$A = \begin{pmatrix} 3 & 2 \\ 1 & 2 \end{pmatrix}, \quad \lambda_1 = 4, \quad \lambda_2 = 1 \quad (\text{eigenvalues})$$

$$x = \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \quad y = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad (\text{eigenvectors})$$



A



$$\alpha x + \beta y \quad \longmapsto \quad \alpha \lambda_1 x + \beta \lambda_2 y$$

Diagonalization:

$$A \in \mathbb{C}^{n \times n} \rightsquigarrow \lambda_1, \lambda_2, \dots, \lambda_n \quad (\text{counted with algebraic multiplicities})$$

$$\rightsquigarrow x^{(1)}, x^{(2)}, \dots, x^{(n)} \quad (\text{associated eigenvectors})$$

$$\rightsquigarrow Ax^{(1)} = \lambda_1 x^{(1)}, \dots, Ax^{(n)} = \lambda_n x^{(n)} \quad (\text{eigenvalue equations})$$

$$A \begin{pmatrix} | & | & \dots & | \\ x^{(1)} & x^{(2)} & \dots & x^{(n)} \\ | & | & \dots & | \end{pmatrix} = \begin{pmatrix} | & | & \dots & | \\ Ax^{(1)} & Ax^{(2)} & \dots & Ax^{(n)} \\ | & | & \dots & | \end{pmatrix}$$

$$= \begin{pmatrix} | & | & \dots & | \\ \lambda_1 x^{(1)} & \lambda_2 x^{(2)} & \dots & \lambda_n x^{(n)} \\ | & | & \dots & | \end{pmatrix} = \underbrace{\begin{pmatrix} | & | & \dots & | \\ x^{(1)} & x^{(2)} & \dots & x^{(n)} \\ | & | & \dots & | \end{pmatrix}}_X \underbrace{\begin{pmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \dots & \\ & & & \lambda_n \end{pmatrix}}_D$$

$$\Rightarrow AX = XD$$

If X is invertible, then: $D = X^{-1}AX$ A is similar to a diagonal matrix

Application:

$$\begin{aligned} A^{98} &= (XD X^{-1})^{98} = XD \underbrace{X^{-1}X}_{\mathbb{1}} D \underbrace{X^{-1}X}_{\mathbb{1}} D X^{-1} \cdots XD X^{-1} \\ &= XD^{98} X^{-1} \\ &= X \begin{pmatrix} \lambda_1^{98} & & \\ & \lambda_2^{98} & \\ & & \ddots \\ & & & \lambda_n^{98} \end{pmatrix} X^{-1} \end{aligned}$$