


Manifolds - Part 5

$$(X, \mathcal{T}) \text{ topological space} \rightsquigarrow (X/\sim, \hat{\mathcal{T}}) \text{ quotient space}$$

Projective space: $P^n(\mathbb{R}) =$ set of 1-dimensional subspaces of \mathbb{R}^{n+1}

$$S^n \subseteq \mathbb{R}^{n+1}$$




$$S^n := \{x \in \mathbb{R}^{n+1} \mid \|x\| = 1\}$$

↖ Euclidean norm


equivalence relation: $x \sim -x$

Let's define: $x \sim y \iff (x=y \text{ or } x=-y)$



$$P^n(\mathbb{R}) := S^n / \sim \text{ with quotient topology}$$

Is $P^n(\mathbb{R})$ a Hausdorff space?




Take $[x]_{\sim}, [y]_{\sim} \in P^n(\mathbb{R})$ with $[x]_{\sim} \neq [y]_{\sim} \implies x \neq y$ and $x \neq -y$

Take open neighbourhoods

$U, V \subseteq S^n$ of x and y , respectively,

with $U \cap V = \emptyset$, $-U \cap V = \emptyset$

$-U \cap -V = \emptyset$, $U \cap -V = \emptyset$



Look at: $\hat{U} := q[U]$, $q: S^n \rightarrow S^n/\sim$ canonical projection

$$q^{-1}[\hat{U}] = U \cup (-U) \xleftarrow{\text{open}} \mathcal{T} \Rightarrow \hat{U} \xleftarrow{\text{open}} \hat{\mathcal{T}}$$

(the same for $\hat{V} := q[V]$)

We find: $q^{-1}[\hat{U} \cap \hat{V}] = q^{-1}[\hat{U}] \cap q^{-1}[\hat{V}] = (U \cup (-U)) \cap (V \cup -V) = \emptyset$

$$\xrightarrow{q \text{ surjective}} \hat{U} \cap \hat{V} = \emptyset$$