

Manifolds - Part 29

M smooth manifold of dimension $n \Rightarrow T_p M$ n -dimensional vector space

Definition:

$$\omega : M \longrightarrow \bigcup_{p \in M} \text{Alt}^k(T_p M)$$

$$p \longmapsto \omega_p = \omega(p) \in \text{Alt}^k(T_p M)$$


is called a k -form on M .

We also define: $\omega \wedge \eta$ as $(\omega \wedge \eta)(p) := \omega(p) \wedge \eta(p)$

$$f^* \omega \quad \text{as} \quad (f^* \omega)(p) := (df_p)^* \omega(p)$$

$$f : N \longrightarrow M \text{ smooth}$$

Basis elements:



basis of $T_p M$: $(\partial_1, \partial_2, \dots, \partial_n)$ with $\partial_j := \varphi_*(e_j) = d\varphi_{h(p)}(e_j)$

basis of $(T_p M)^* = \text{Alt}^1(T_p M)$: $(dx_p^1, dx_p^2, \dots, dx_p^n)$

$$\text{defined by: } dx_p^j(\partial_k) = \delta_k^j = \begin{cases} 1, & j=k \\ 0, & j \neq k \end{cases}$$

Proposition: A basis of $\text{Alt}^k(T_p M)$ is given by:

$$(dx_p^{\mu_1} \wedge dx_p^{\mu_2} \wedge \dots \wedge dx_p^{\mu_k})_{\mu_1 < \mu_2 < \dots < \mu_k}$$

Example: $\dim(M) = 3$, $\text{Alt}^2(\mathcal{T}_p M)$:

$$(dx_p^1 \wedge dx_p^2, dx_p^1 \wedge dx_p^3, dx_p^2 \wedge dx_p^3)$$

Conclusion: Each k -form on M can locally be written as:

$$\omega(p) = \sum_{\mu_1 < \dots < \mu_k} \omega_{\mu_1, \mu_2, \dots, \mu_k}(p) \cdot dx_p^{\mu_1} \wedge dx_p^{\mu_2} \wedge \dots \wedge dx_p^{\mu_k}$$

$$\omega_{\mu_1, \mu_2, \dots, \mu_k} : U \longrightarrow \mathbb{R} \quad \text{component functions}$$

Definition: • If all component functions are differentiable at p ,
then ω is differentiable at p .

• If ω is differentiable at all $p \in M$,

then ω is called a differential form on M .

$$\omega \in \Omega^k(M)$$

$$\Omega^0(M) := C^\infty(M)$$