



Manifolds - Part 41

We already know:

$$\int_{\mathsf{A}} \omega := \int_{\mathsf{h}[\mathsf{A}]} \varphi^* \omega$$



 ω canonical volume form on S^{1} (measures areas on S^{2}) Example:



$$\int \omega = \int_{\widetilde{\mathbf{u}}} \Phi^* \omega$$

$$\underline{\Phi}[\widetilde{\mathbf{u}}]$$

canonical volume form: $\omega(p) = \sqrt{\det(G(p))} dx_p^1 \wedge dx_p^2$ $\sin(\theta) \qquad d\theta \qquad d\phi$ for $p = \Phi(\theta, \phi)$ 1-forms on S^2

$$\left(\underbrace{\Phi}^* \omega \right) \left(\underbrace{\tilde{p}} \right) = \sin(\theta) \cdot \det(\cdot, \cdot)$$

$$d\theta \wedge d\phi$$

$$1 - \text{forms on } \widetilde{\mathbb{N}} \subseteq \mathbb{R}^2$$

in short:
$$\omega = \sin(\theta) d\theta \wedge d\phi$$

$$\Phi^* \omega = \sin(\theta) d\theta \wedge d\phi$$


$$\int_{S^{1}\setminus\{\dots\}} \omega = \int_{S^{1}} \Phi[\tilde{u}] = \int_{(0,\pi)\times(0,2\pi)}^{*} \omega = \int_{(0,\pi)\times(0,2\pi)}^{*} \sin(\theta) d\theta \wedge d\phi$$

$$= \int_{0}^{*} \left(\int_{0}^{*} \sin(\theta) d\phi\right) d\theta = 4\pi$$

Definition: Let M be an orientable
$$n$$
-dimensional manifold and $\omega \in \Omega^n(M)$.

A set $A \subseteq M$ is called

measurable if h[An U] is measurable
 (w.r.t. Lebesque measure in Rⁿ)



• <u>null set</u> (set with measure zero) if h[AnU] has Lebesgue measure 0 for every chart (U,h).

We get: $\int_{A} \omega \quad \text{is defined for every measurable set } A \subseteq U \quad \text{(where (u,h) is a chart)}$ $\text{(assuming } \int_{h[A]} \psi^* \omega \quad \text{exists in } \mathbb{R}^n \text{)}$ $\text{and } \int_{A} \omega := \int_{A} \omega \quad \text{if } B \setminus N \subseteq U \quad \text{(where (u,h) is a chart)}$

$$\frac{\text{Hence:}}{S^{2}} \qquad \int_{S^{2}} \omega = 4 \, \text{m}$$