




# Manifolds - Part 43



Proposition: (well-definedness of  $\int_A \omega$ )


$(U_k, h_k)_{k \in \mathbb{N}}$  atlas,  $A = \bigcup_{k \in \mathbb{N}} A_k$  disjoint  $A_k \subseteq U_k$  with:

(1)  $\int_{A_k} \omega$  exists for all  $k \in \mathbb{N}$



(2)  $\sum_{k=1}^{\infty} \int_{h_k[A_k]} |\omega_{1,2,\dots,n}(h_k^{-1}(x))| d^n x < \infty$

$(\tilde{U}_m, \tilde{h}_m)_{m \in \mathbb{N}}$  atlas,  $A = \bigcup_{m \in \mathbb{N}} \tilde{A}_m$  disjoint  $\tilde{A}_m \subseteq \tilde{U}_m$  (measurable).




Then:

(1-tilde)  $\int_{\tilde{A}_m} \omega$  exists for all  $m \in \mathbb{N}$

(2-tilde)  $\sum_{m=1}^{\infty} \int_{\tilde{h}_m[\tilde{A}_m]} |\omega_{1,2,\dots,n}(\tilde{h}_m^{-1}(x))| d^n x < \infty$

and:

$$\sum_{m=1}^{\infty} \int_{\tilde{h}_m[\tilde{A}_m]} \omega_{1,2,\dots,n}(\tilde{h}_m^{-1}(x)) d^n x = \sum_{k=1}^{\infty} \int_{h_k[A_k]} \omega_{1,2,\dots,n}(h_k^{-1}(x)) d^n x = \int_A \omega$$



part 40

$$\int_{h_k[A_k \cap \tilde{A}_m]} |\omega_{1,2,\dots,n}(h_k^{-1}(x))| d^n x = \int_{\tilde{h}_m[A_k \cap \tilde{A}_m]} |\omega_{1,2,\dots,n}(\tilde{h}_m^{-1}(x))| d^n x$$

$$\Rightarrow \sum_{m=1}^{\infty} \int_{h_k[A_k \cap \tilde{A}_m]} |\omega_{1,2,\dots,n}(h_k^{-1}(x))| d^n x = \sum_{m=1}^{\infty} \int_{\tilde{h}_m[A_k \cap \tilde{A}_m]} |\omega_{1,2,\dots,n}(\tilde{h}_m^{-1}(x))| d^n x$$

$$\equiv \int_{\bigcup_{m \in \mathbb{N}} h_k[A_k \cap \tilde{A}_m]} |\omega_{1,2,\dots,n}(h_k^{-1}(x))| d^n x$$

$$\equiv \int_{h_k[A_k]} |\omega_{1,2,\dots,n}(h_k^{-1}(x))| d^n x$$

finite!

$$\Rightarrow \sum_{k=1}^{\infty} \int_{h_k[A_k]} |\omega_{1,2,\dots,n}(h_k^{-1}(x))| d^n x = \sum_{k=1}^{\infty} \sum_{m=1}^{\infty} \int_{\tilde{h}_m[A_k \cap \tilde{A}_m]} |\omega_{1,2,\dots,n}(\tilde{h}_m^{-1}(x))| d^n x$$

$$= \sum_{m=1}^{\infty} \int_{\tilde{h}_m[\tilde{A}_m]} |\omega_{1,2,\dots,n}(\tilde{h}_m^{-1}(x))| d^n x$$

same calculation without absolute value

$$\Rightarrow \sum_{k=1}^{\infty} \int_{h_k[A_k]} \omega_{1,2,\dots,n}(h_k^{-1}(x)) d^n x = \sum_{m=1}^{\infty} \int_{\tilde{h}_m[\tilde{A}_m]} \omega_{1,2,\dots,n}(\tilde{h}_m^{-1}(x)) d^n x$$

□