


## **The Bright Side of Mathematics**

The following pages cover the whole Manifolds course of the Bright Side of Mathematics. Please note that the creator lives from generous supporters and would be very happy about a donation. See more here: <https://tbsom.de/support>


Have fun learning mathematics!

# Manifolds - Part 1

generalised surfaces?




How to calculate on them?



⇒ (generalised) Stokes's Theorem

Metric space:

$(X, d)$   
 ↑ set      ↑ distance function



↷ define open sets  $A \subseteq X$

Definition:

Let  $X$  be a set,  $\mathcal{P}(X)$  be the power set,  
 and  $\mathcal{T} \subseteq \mathcal{P}(X)$  be a collection of subsets.

If  $\mathcal{T}$  satisfies: (1)  $\emptyset, X \in \mathcal{T}$

(2)  $A, B \in \mathcal{T} \Rightarrow A \cap B \in \mathcal{T}$

(3)  $(A_i)_{i \in I}$  with  $A_i \in \mathcal{T} \Rightarrow \bigcup_{i \in I} A_i \in \mathcal{T}$

then  $\mathcal{T}$  is called a topology on  $X$ .

The elements of  $\mathcal{T}$  are called open sets.

Examples:

(a)  $\mathcal{T} = \{\emptyset, X\}$  is a topology on  $X$  (indiscrete topology)





(b)  $\mathcal{T} = \mathcal{P}(X)$  is a topology on  $X$  (discrete topology)

# Manifolds - Part 2

- $\mathcal{T} \subseteq \mathcal{P}(X)$  topology on  $X$ :
- (1)  $\emptyset, X \in \mathcal{T}$
  - (2)  $A, B \in \mathcal{T} \Rightarrow A \cap B \in \mathcal{T}$
  - (3)  $(A_i)_{i \in I}$  with  $A_i \in \mathcal{T} \Rightarrow \bigcup_{i \in I} A_i \in \mathcal{T}$

$(X, \mathcal{T})$  is called a topological space.

Important names:  $(X, \mathcal{T})$  topological space,  $S \subseteq X$ ,  $p \in X$

- (a)  $p$  interior point of  $S$   $:\Leftrightarrow$  There is an open set  $U \in \mathcal{T}$ :  
 $p \in U$  and  $U \subseteq S$  
- (b)  $p$  exterior point of  $S$   $:\Leftrightarrow$  There is an open set  $U \in \mathcal{T}$ :  
 $p \in U$  and  $U \subseteq X \setminus S$  
- (c)  $p$  boundary point of  $S$   $:\Leftrightarrow$  For all open sets  $U \in \mathcal{T}$  with  $p \in U$ :  
 $U \cap S \neq \emptyset$  and  $U \cap (X \setminus S) \neq \emptyset$  
- (d)  $p$  accumulation point of  $S$   $:\Leftrightarrow$  For all open sets  $U \in \mathcal{T}$  with  $p \in U$ :  
 $U \setminus \{p\} \cap S \neq \emptyset$  

- More names:
- (a)  $S^\circ := \{p \in X \mid p \text{ interior point of } S\}$  interior of  $S$
  - (b)  $\text{Ext}(S) := \{p \in X \mid p \text{ exterior point of } S\}$  exterior of  $S$
  - (c)  $\partial S := \{p \in X \mid p \text{ boundary point of } S\}$  boundary of  $S$
  - (d)  $S' := \{p \in X \mid p \text{ accumulation point of } S\}$  derived set of  $S$
  - (e)  $\bar{S} := S \cup \partial S$  closure of  $S$

Example:  $X = \mathbb{R}$ ,  $\mathcal{T} = \{\emptyset, \mathbb{R}\} \cup \{(a, \infty) \mid a \in \mathbb{R}\}$

$S = (0, 1)$  ← not an open set!

← no interior points: there is no  $\emptyset \neq U \in \mathcal{T}$  with  $U \subseteq S$


$$\Rightarrow S^\circ = \emptyset$$

$$X \setminus S = (-\infty, 0] \cup [1, \infty) \Rightarrow \text{Ext}(S) = (1, \infty)$$


$$\Rightarrow \partial S = (-\infty, 1] \Rightarrow \bar{S} = (-\infty, 1]$$

# Manifolds - Part 3


$(X, \mathcal{T})$  topological space



Convergence:  $(a_n)_{n \in \mathbb{N}}$ ,  $a_n \in X$   
converges to  $a \in X$

In a metric space:  The sequence members lie in each  $\epsilon$ -ball around  $a$ , eventually.

For each  $\epsilon$ -ball  $B_\epsilon(a)$ , there is  $N \in \mathbb{N}$  such that  
for all  $n \geq N$ :  $a_n \in B_\epsilon(a)$

In a topological space:    
open neighbourhood of  $a$   
an open set  $U \in \mathcal{T}$  with  $a \in U$

Definition:  $(X, \mathcal{T})$  topological space,  $(a_n)_{n \in \mathbb{N}}$  sequence in  $X$ .

$a_n \xrightarrow{n \rightarrow \infty} a \iff$  For each  $U \in \mathcal{T}$  with  $a \in U$ , there is  $N \in \mathbb{N}$   
such that for all  $n \geq N$ :  $a_n \in U$

Example:  $X = \mathbb{R}$ ,  $\mathcal{T} = \{\emptyset, \mathbb{R}\} \cup \{(b, \infty) \mid b \in \mathbb{R}\}$


$$(a_n)_{n \in \mathbb{N}} = \left(\frac{1}{n}\right)_{n \in \mathbb{N}}$$

- converges to  $0$ : each open neighbourhood of  $0$  looks like  $(b, \infty)$  for  $b < 0$ , so:  $\frac{1}{n} \in (b, \infty)$
- converges to  $-1$ : each open neighbourhood of  $-1$  looks like  $(b, \infty)$  for  $b < -1$ , so:  $\frac{1}{n} \in (b, \infty)$
- converges to  $-2$

Definition: A topological space  $(X, \mathcal{T})$  is called a Hausdorff space if

for all  $x, y \in X$  with  $x \neq y$  there is an open neighbourhood of  $x$ :  $U_x \in \mathcal{T}$


and there is an open neighbourhood of  $y$ :  $U_y \in \mathcal{T}$



with:  $U_x \cap U_y = \emptyset$

# Manifolds - Part 4

Projective space:  $P^n(\mathbb{R}) =$  set of 1-dimensional subspaces of  $\mathbb{R}^{n+1}$



the directions define a set + topology?


Quotient topology:  $(X, \mathcal{T})$  topological space,  $\sim$  equivalence relation on  $X$

- $\hookrightarrow$  reflexive  $x \sim x$
- symmetric  $x \sim y \Rightarrow y \sim x$
- transitive  $x \sim y \wedge y \sim z \Rightarrow x \sim z$

equivalence class of  $x$  :  $[x]_{\sim} := \{y \in X \mid y \sim x\}$

$X/\sim := \{[x]_{\sim} \mid x \in X\}$  quotient set

$q: X \rightarrow X/\sim, x \mapsto [x]_{\sim}$  canonical projection





$q^{-1}[U] \subseteq X$  open  $\iff U \subseteq X/\sim$  open

$q^{-1}[U] \in \mathcal{T}$   $\iff U \in \hat{\mathcal{T}}$

This defines a topology  $\hat{\mathcal{T}}$  on  $X/\sim$ , called the quotient topology.

Example:

$X = [0,1] \times (-1,1)$




equivalence relation:  $(0,s) \sim (1,-s)$

# Manifolds - Part 5

$$(X, \mathcal{T}) \text{ topological space} \rightsquigarrow (X/\sim, \hat{\mathcal{T}}) \text{ quotient space}$$

Projective space:  $P^n(\mathbb{R}) = \text{set of 1-dimensional subspaces of } \mathbb{R}^{n+1}$

$$S^n \subseteq \mathbb{R}^{n+1}$$




$$S^n := \{x \in \mathbb{R}^{n+1} \mid \|x\| = 1\}$$

↖ Euclidean norm


equivalence relation:  $x \sim -x$

Let's define:  $x \sim y \iff (x=y \text{ or } x=-y)$



$$P^n(\mathbb{R}) := S^n / \sim \text{ with quotient topology}$$

Is  $P^n(\mathbb{R})$  a Hausdorff space?




Take  $[x]_{\sim}, [y]_{\sim} \in P^n(\mathbb{R})$  with  $[x]_{\sim} \neq [y]_{\sim} \implies x \neq y$  and  $x \neq -y$

Take open neighbourhoods

$U, V \subseteq S^n$  of  $x$  and  $y$ , respectively,

with  $U \cap V = \emptyset$ ,  $-U \cap V = \emptyset$

$-U \cap -V = \emptyset$ ,  $U \cap -V = \emptyset$





Look at:  $\hat{u} := q[u]$ ,  $q: S^n \rightarrow S^n / \sim$  canonical projection

$$q^{-1}[\hat{u}] = \cup (-u) \xleftarrow{\text{open}} \mathcal{J} \Rightarrow \hat{u} \xleftarrow{\text{open}} \hat{\mathcal{J}}$$

(the same for  $\hat{v} := q[v]$ )

We find:  $q^{-1}[\hat{u} \cap \hat{v}] = q^{-1}[\hat{u}] \cap q^{-1}[\hat{v}] = (\cup (-u)) \cap (\cup (-v)) = \emptyset$

$$\xrightarrow{q \text{ surjective}} \hat{u} \cap \hat{v} = \emptyset$$

## Manifolds - Part 6


$(X, \mathcal{T})$  topological space: generate the topology  $\mathcal{T}$

Definition: Let  $(X, \mathcal{T})$  be a topological space. A collection of open subsets

$\mathcal{B} \subseteq \mathcal{T}$  is called a basis (base) of  $\mathcal{T}$  if:

for all  $U \in \mathcal{T}$  there is  $(A_i)_{i \in I}$  with  $A_i \in \mathcal{B}$

and  $\bigcup_{i \in I} A_i = U$



Examples: (a)  $\mathcal{B} = \mathcal{T}$  is always a basis.

(b) If  $\mathcal{T}$  is discrete topology on  $X$ , then  $\mathcal{B} = \{\{x\} \mid x \in X\}$  is a basis of  $\mathcal{T}$ .

(c) Let  $(X, \mathcal{T})$  be the topological space induced by a metric space  $(X, d)$   
 $\mathcal{B} = \{B_\epsilon(x) \mid x \in X, \epsilon > 0\}$  is a basis of  $\mathcal{T}$ .




(d)  $\mathbb{R}^n$  with standard topology (defined by Euclidean metric)

$\mathcal{B} = \{B_\epsilon(x) \mid x \in \mathbb{Q}^n, \epsilon \in \mathbb{Q}, \epsilon > 0\}$  is a basis of  $\mathcal{T}$ .

*only countably many elements*

Definition: A topological space  $(X, \mathcal{T})$  is called second-countable if there is a countable basis of  $\mathcal{T}$ .


# Manifolds - Part 7



Definition:  $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$  topological spaces.

$f: X \rightarrow Y$  is called continuous if

$$U \in \mathcal{T}_Y \Rightarrow f^{-1}[U] \in \mathcal{T}_X.$$



homeomorphism =  $f: X \rightarrow Y$  bijective, continuous and  $f^{-1}: Y \rightarrow X$  continuous

Examples: (a)  $(Y, \mathcal{T}_Y) =$  indiscrete topological space  $\Rightarrow f: X \rightarrow Y$  continuous

(b)  $(X, \mathcal{T}_X) =$  discrete topological space  $\Rightarrow f: X \rightarrow Y$  continuous

(c)  $(X, \mathcal{T}_X)$  with equivalence relation  $\sim$ ,  $(X/\sim, \hat{\mathcal{T}})$  quotient space

$q: X \rightarrow X/\sim, x \mapsto [x]_{\sim}$  canonical projection  
is continuous


Definition:  $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$  topological spaces.

$f: X \rightarrow Y$  is called sequentially continuous if for all  $x \in X$ :

$(x_n)_{n \in \mathbb{N}} \subseteq X$  with  $x_n \xrightarrow{n \rightarrow \infty} x$

$\Rightarrow$

$(f(x_n))_{n \in \mathbb{N}} \subseteq Y$  convergent with  $f(x_n) \xrightarrow{n \rightarrow \infty} f(x)$



Fact:

$f: X \rightarrow Y$  continuous  $\iff$   $f: X \rightarrow Y$  sequentially continuous


in metric spaces

second-countable spaces

# Manifolds - Part 8

$[a, b] \subseteq \mathbb{R}$  compact (Bolzano-Weierstrass and Heine-Borel)

$(X, \mathcal{T})$




cover with open sets  
 $\Downarrow$   
 do finitely many suffice?

Definition: Let  $(X, \mathcal{T})$  be a topological space and  $A \subseteq X$ .

$A$  is called compact if

$\bigcup_{i \in I} U_i \supseteq A$  with  $U_i \in \mathcal{T} \Rightarrow$  there is a finite  $I_0 \subseteq I$  with:  $\bigcup_{i \in I_0} U_i \supseteq A$




We know:  $A \subseteq \mathbb{R}^n$  compact  $\Leftrightarrow A$  closed and bounded (Heine-Borel theorem)  
with standard topology

Proposition: Let  $(X, \mathcal{T})$  be a Hausdorff space. Then:

$A \subseteq X$  compact  $\Rightarrow A$  closed  $\left( \begin{array}{l} X \setminus A \text{ open} \\ X \setminus A \in \mathcal{T} \end{array} \right)$



Proof:



Assume  $A$  is compact.

Fix  $b \in X \setminus A$ .

For any  $a \in A$ , there are  $U_a, V_a \in \mathcal{T}$   
with  $a \in U_a$ ,  $b \in V_a$  and  $U_a \cap V_a = \emptyset$




$$A \subseteq \bigcup_{a \in A} U_a \quad (\text{open cover})$$

$A$  compact

$$\Rightarrow A \subseteq \bigcup_{j=1}^n U_{a_j} \quad (\text{finite subcover})$$

$$\Rightarrow V := \bigcap_{j=1}^n V_{a_j} \quad \text{open neighbourhood of } b$$




$$\text{with } A \cap V \subseteq \bigcup_{j=1}^n U_{a_j} \cap \bigcap_{j=1}^n V_{a_j} = \emptyset$$

$$\Rightarrow b \text{ is an interior point of } X \setminus A \Rightarrow A \text{ closed}$$

# Manifolds - Part 9

Definition:  $n$ -dimensional (topological) manifold:


- topological space  $(X, \mathcal{T})$  with:
- (1) Hausdorff space
  - (2) second-countable
  - (3) locally Euclidean of dimension  $n$




Definition:  $(X, \mathcal{T})$  is called locally Euclidean of dimension  $n$  if:

For all  $x \in X$  there is an open neighbourhood  $U \in \mathcal{T}$  and a homeomorphism  $h: U \rightarrow U'$  with  $U' \subseteq \mathbb{R}^n$  open.

The map  $h: U \rightarrow U'$  is called a chart of  $(X, \mathcal{T})$ .



# Manifolds - Part 10



(1) Hausdorff space


(2) second-countable

(3) locally Euclidean of dimension  $n$


chart  $(U, h)$

Definition: A collection of charts  $(U_i, h_i)_{i \in I}$  is called an atlas if:  $\bigcup_{i \in I} U_i = M$

Example: (a)  $(M, \mathcal{T})$  discrete topological space with countably many points



(b)  $M \subseteq \mathbb{R}^n$  open subset,  $(M, \mathcal{T})$  with standard topology




(c)  $S^2 \subseteq \mathbb{R}^3$ ,  $S^2 := \{x \in \mathbb{R}^3 \mid \|x\| = 1\}$

2-dimensional manifold

$$h_{3,-}: \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \mapsto \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

$$h_{3,-}^{-1}: \begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} \mapsto \begin{pmatrix} x'_1 \\ x'_2 \\ -\sqrt{1 - \|x'\|^2} \end{pmatrix}$$



$(U_{i,\pm}, h_{i,\pm})_{i \in \{1,2,3\}}$  is an atlas.



# Manifolds - Part 11

$$S^n := \{x \in \mathbb{R}^{n+1} \mid \|x\| = 1\}$$




$$= \{x \in \mathbb{R}^{n+1} \mid \pm x_i > 0\}$$

is an  $n$ -dimensional manifold with atlas  $(U_{i,\pm}, h_{i,\pm})_{i \in \{1, \dots, n+1\}}$

Projective space:  $P^n(\mathbb{R}) := S^n / \sim$  with quotient topology

equivalence relation:  $x \sim y \Leftrightarrow (x=y \text{ or } x=-y)$



$$q: S^n \rightarrow S^n / \sim \quad \text{canonical projection}$$


$$x \mapsto [x]_{\sim}$$

$$V_i := \{[x]_{\sim} \in P^n(\mathbb{R}) \mid x_i \neq 0\}, \quad q^{-1}[V_i] = U_{i,+} \cup U_{i,-}$$

$\hookrightarrow$  open

for  $n=1$ :  $h_1: V_1 \rightarrow V_1' \subseteq \mathbb{R}^1, \quad h_1([x]_{\sim}) = \frac{x_2}{x_1}$  slope

with inverse  $h_1^{-1}(x_1') = \left[ \begin{pmatrix} 1 \\ x_1' \end{pmatrix} \cdot \frac{1}{\sqrt{1+(x_1')^2}} \right]_{\sim}$




$h_2$  works similarly  $\Rightarrow$  1-dimensional manifold

for  $n \in \mathbb{N}$ :  $h_i: V_i \rightarrow V_i' \subseteq \mathbb{R}^n$

$$h_i([x]_{\sim}) = \begin{pmatrix} \frac{x_1}{x_i} \\ \vdots \\ \frac{x_{i-1}}{x_i} \\ \frac{x_{i+1}}{x_i} \\ \vdots \\ \frac{x_{n+1}}{x_i} \end{pmatrix} \quad \text{homeomorphism}$$

$\Rightarrow$   $n$ -dimensional manifold

# Manifolds - Part 12



- $C^k$ -diffeomorphism:
- $k \in \{0, 1, \dots\}$
  - or  $k = \infty$
- $\omega$  is  $k$ -times continuously differentiable (partial derivatives up to the  $k$ -th order exist and are continuous)
  - $\omega$  is bijective
  - $\omega^{-1} \in C^k(\dots)$
- }  $\omega \in C^k(\cdot)$

Definition: • Two charts  $h, k$  are called  $C^k$ -smoothly compatible if the transition map is a  $C^k$ -diffeomorphism.

- An atlas  $\{(U_i, h_i)_{i \in I}\}$  is called a  $C^k$ -atlas if any two charts are  $C^k$ -smoothly compatible.
- A maximal  $C^k$ -atlas  $\mathcal{A}$  is:
  - (1)  $\mathcal{A}$  is a  $C^k$ -atlas
  - (2) For any other  $C^k$ -atlas  $\mathcal{B}$ , we have  $\mathcal{B} \not\supseteq \mathcal{A}$ .

Definition:  $n$ -dimensional  $C^k$ -smooth manifold:

- $n$ -dimensional (topological) manifold
- maximal  $C^k$ -atlas ( $C^k$ -smooth structure)


# Manifolds - Part 13

Examples for smooth manifolds:


(a)  $S^n \subseteq \mathbb{R}^{n+1}$  is a smooth manifold.

We show that  $(U_{i,\pm}, h_{i,\pm})_{i \in \{1, \dots, n+1\}}$  is  $C^\infty$ -atlas:

$$\{x \in \mathbb{R}^{n+1} \mid \pm x_i > 0\}$$



$$h_{i,\pm} : \begin{pmatrix} x_1 \\ \vdots \\ x_{i-1} \\ x_{i+1} \\ \vdots \\ x_{n+1} \end{pmatrix} \mapsto \begin{pmatrix} x_1 \\ \vdots \\ x_{i-1} \\ x_{i+1} \\ \vdots \\ x_{n+1} \end{pmatrix}$$



For  $n=2, i=3, j=1$


$$x' = \begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} \xrightarrow{h_{i,+}^{-1}} \begin{pmatrix} x'_1 \\ x'_2 \\ \sqrt{1 - \|x'\|^2} \end{pmatrix} \xrightarrow{h_{j,+}} \begin{pmatrix} x'_2 \\ \sqrt{1 - \|x'\|^2} \end{pmatrix} \quad C^\infty\text{-diffeomorphism}$$

$\rightsquigarrow$  extend to a maximal  $C^\infty$ -atlas  $\rightsquigarrow$   $C^\infty$ -smooth manifold

(b)  $\mathbb{R}^n$  is a smooth manifold

$\hookrightarrow$  atlas given by one chart  $(\mathbb{R}^n, id)$   $\rightsquigarrow$  extend to a maximal  $C^\infty$ -atlas  
(standard smooth structure for  $\mathbb{R}^n$ )

(c) Consider  $f \in C^1(\mathbb{R})$




$$G_f = \{(x, f(x)) \mid x \in \mathbb{R}\} \\ \subseteq \mathbb{R} \times \mathbb{R}$$

$G_f$  is a 1-dimensional manifold with one chart:  $h: G_f \rightarrow \mathbb{R}$

$$(x, f(x)) \mapsto x$$

$\rightsquigarrow$  extend to a smooth structure


# Manifolds - Part 14



Definition: Let  $M$  be an  $n$ -dimensional (smooth) manifold.  
 $M_0 \subseteq M$  is called a  $k$ -dimensional submanifold of  $M$  if

for all  $p \in M_0$  there is a chart  $(u, h)$  of  $M$  with

$$h[M_0 \cap U] = (\mathbb{R}^k \times \underbrace{0}_{n-k \text{ zeros}}) \cap U'$$



$(u, h)$  is called a submanifold chart for  $M_0$ .

Note:  $M_0$  is also a manifold:


$(u, h)$  submanifold chart  $\rightsquigarrow (\tilde{u}, \tilde{h})$  chart,  $\tilde{u} := U \cap M_0$

$$\tilde{h} \text{ given by } p \mapsto h(p) = \begin{pmatrix} \otimes \\ \vdots \\ \otimes \\ 0 \\ \vdots \\ 0 \end{pmatrix} \mapsto \begin{pmatrix} \otimes \\ \vdots \\ \otimes \end{pmatrix} \in \mathbb{R}^k$$

## Manifolds - Part 15

Regular value theorem in  $\mathbb{R}^n$  = preimage theorem = submersion theorem

$$f: \mathbb{R}^n \rightarrow \mathbb{R}^m \text{ smooth}$$



preimage = smooth submanifold?

Definition:  $f: U \rightarrow \mathbb{R}^m$ ,  $U \subseteq \mathbb{R}^n$  open,  $C^1$ -function.

(1)  $x \in U$  is called a critical point of  $f$  if  $df_x$  is not surjective (or  $J_f(x)$  has rank less than  $m$ )

(2)  $c \in f[U]$  is called a regular value of  $f$  if  $f^{-1}[\{c\}]$  does not contain any critical points.

Theorem:

$$f: U \rightarrow \mathbb{R}^m, U \subseteq \mathbb{R}^n \text{ open, } C^\infty\text{-function. } (n \geq m)$$

If  $c$  is a regular value of  $f$ , then

$f^{-1}[\{c\}]$  is an  $(n-m)$ -dimensional submanifold of  $\mathbb{R}^n$ .

Proof: Use implicit function theorem.

Example:

$$f: \mathbb{R}^n \rightarrow \mathbb{R}, f(x_1, \dots, x_n) = x_1^2 + x_2^2 + \dots + x_n^2$$

$$J_f(x_1, \dots, x_n) = (2x_1 \quad 2x_2 \quad \dots \quad 2x_n)$$

$\Rightarrow x=0$  is the only critical point.


Hence:  $1$  is a regular value.

$$\Rightarrow f^{-1}[\{1\}] = S^{n-1} \text{ submanifold of } \mathbb{R}^n.$$

# Manifolds - Part 16

Smooth maps:

Use the smooth structures!



Definition: Let  $M$  and  $N$  be  $C^\infty$ -smooth manifolds.

A map  $f: M \rightarrow N$  is called  $k$ -times differentiable at  $p \in M$


if for charts  $(U, h), (W, k)$  with  $p \in U$  and  $f(p) \in W$

the map  $k \circ f \circ h^{-1}$   $k$ -times differentiable at  $h(p)$ .

Moreover:  $f: M \rightarrow N$  is called  $C^\infty$ -smooth if  $f$  is  $k$ -times differentiable at  $p \in M$

for every  $p \in M$  and every  $k \in \mathbb{N}$ . We write:  $f \in C^\infty(M, N)$ .

# Manifolds - Part 17



Examples of smooth maps:

(1)  $S^2 \longrightarrow \mathbb{R}^3$


inclusion map:

$i: X \mapsto X$

continuous!  
smooth?

$$h_{3,-} \left( \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \right) = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$


$$h_{3,-}^{-1} \left( \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) = \begin{pmatrix} x_1 \\ x_2 \\ -\sqrt{1 - \|x'\|^2} \end{pmatrix}$$



$k \circ i \circ h_{3,-}^{-1} : \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \mapsto \begin{pmatrix} x_1 \\ x_2 \\ -\sqrt{1 - \|x'\|^2} \end{pmatrix}$  differentiable  $\implies i$  is smooth




(2)  $q: S^2 \rightarrow P^2(\mathbb{R}) = S^2/\sim$  ( $x \sim y \Leftrightarrow x = y$  or  $x = -y$ )  
 $x \mapsto [x]_{\sim}$  continuous map! smooth?



# Manifolds - Part 18

Regular Value Theorem:




Let  $M, N$  be smooth manifolds of dimension  $m$  and  $n$  ( $m \geq n$ ),  
 $f: M \rightarrow N$  be a smooth map, and  $q \in N$  be a regular value of  $f$ .

↳  $f^{-1}[\{q\}]$  does not contain critical points

↳  $p \in M$  is called a critical point of  $f$  if  
 $\text{rank } f_p := \text{rank} \left( J_{k \circ f \circ h^{-1}}(h(p)) \right)$   
 is less than  $n$  (not maximal!).

Then:  $f^{-1}[\{q\}]$  is a  $(m-n)$ -dim submanifold of  $M$ .

Example: (a)  $GL(d, \mathbb{R}) := \{A \in \mathbb{R}^{d \times d} \mid \det(A) \neq 0\}$  is manifold of dimension  $d^2$ .

(b)  $\text{Sym}(d \times d, \mathbb{R}) := \{B \in \mathbb{R}^{d \times d} \mid B^T = B\}$  is manifold of dimension  $\frac{d(d+1)}{2}$   
 $\frac{d^2-d}{2}$    $d^2 - \frac{d^2-d}{2} = \frac{d(d+1)}{2}$


(c)  $O(d, \mathbb{R}) := \{A \in GL(d, \mathbb{R}) \mid A^T A = \mathbb{1}\}$  is a submanifold of  $GL(d, \mathbb{R})$

Proof:  $f: GL(d, \mathbb{R}) \longrightarrow \text{Sym}(d \times d, \mathbb{R})$  ,  $f(A) = A^T A$

Two things to show: (1)  $f^{-1}[\{\mathbb{1}\}] = O(d, \mathbb{R})$

(2)  $\mathbb{1}$  is a regular value of  $f$

Case  $d=2$ :




$$\begin{aligned} (k \circ f \circ h^{-1}) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} &= (k \circ f) \begin{pmatrix} x_1 & x_2 \\ x_3 & x_4 \end{pmatrix} = k \left( \begin{pmatrix} x_1 & x_2 \\ x_3 & x_4 \end{pmatrix}^T \begin{pmatrix} x_1 & x_2 \\ x_3 & x_4 \end{pmatrix} \right) \\ &= k \left( \begin{pmatrix} x_1^2 + x_3^2 & x_1 x_2 + x_3 x_4 \\ x_1 x_2 + x_3 x_4 & x_2^2 + x_4^2 \end{pmatrix} \right) = \begin{pmatrix} x_1^2 + x_3^2 \\ x_1 x_2 + x_3 x_4 \\ x_2^2 + x_4^2 \end{pmatrix} \end{aligned}$$

Jacobian matrix:  $J_{k \circ f \circ h^{-1}}(x) = \begin{pmatrix} 2x_1 & 0 & 2x_3 & 0 \\ x_2 & x_1 & x_4 & x_3 \\ 0 & 2x_2 & 0 & 2x_4 \end{pmatrix}$

rank = 3? Not for:  $x_1 = x_2 = 0$   
 $x_3 = x_4 = 0$   
 $x_1 = x_3 = 0$   
 $x_2 = x_4 = 0$

If  $f(A) = \mathbb{1} \implies J_{k \circ f \circ h^{-1}}(h(A))$  has rank 3  $\implies \mathbb{1}$  regular value


$\implies O(d, \mathbb{R})$  is a submanifold of dimension  $d^2 - \frac{d(d+1)}{2} = \frac{d(d-1)}{2}$




## Manifolds - Part 19

submanifold:  $M \subseteq \mathbb{R}^n$   $k$ -dimensional submanifold

$$h[M \cap U] = (\mathbb{R}^k \times \underbrace{0}_{n-k \text{ zeros}}) \cap U'$$




Example:




$$\begin{aligned} \varphi: \mathbb{R}^1 \cap U' &\rightarrow M \cap U \\ t &\mapsto \begin{pmatrix} \cos(t) \\ \sin(t) \end{pmatrix} \end{aligned}$$

Tangent space:



$$\begin{aligned} T_p^{\text{sub}} M &:= d\varphi_{\tilde{p}}[\mathbb{R}^k] \\ &= \left\{ J_\varphi(\tilde{p})x \mid x \in \mathbb{R}^k \right\} \subseteq \mathbb{R}^n \end{aligned}$$

Example:



surface given by a graph of a function:

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f \in C^1(\mathbb{R}^2)$$

$$M = G_f := \left\{ \begin{pmatrix} x \\ y \\ f(x,y) \end{pmatrix} \mid (x,y) \in \mathbb{R}^2 \right\}$$

parameterisation:  $\varphi: \mathbb{R}^2 \rightarrow M$ ,  $\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ y \\ f(x,y) \end{pmatrix}$


$$J_{\varphi}(x,y) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ \frac{\partial f}{\partial x}(x,y) & \frac{\partial f}{\partial y}(x,y) \end{pmatrix}$$

$$\Rightarrow T_p^{\text{sub}} M = \text{span} \left( \begin{pmatrix} 1 \\ 0 \\ \frac{\partial f}{\partial x}(x,y) \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ \frac{\partial f}{\partial y}(x,y) \end{pmatrix} \right)$$

$p = \begin{pmatrix} x \\ y \\ f(x,y) \end{pmatrix}$

# Manifolds - Part 20

$T_p^{\text{sub}} M$  tangent space for submanifold  $M \subseteq \mathbb{R}^n$ ,  $p \in M$



$$T_p^{\text{sub}} M := \left\{ J_\psi(\tilde{\varphi}^{-1}(p)) x \mid x \in \mathbb{R}^k \right\} \subseteq \mathbb{R}^n$$


Idea:



Proposition:  $T_p^{\text{sub}} M = \left\{ \gamma'(0) \mid \gamma: (-\epsilon, \epsilon) \rightarrow M \text{ differentiable with } \gamma(0) = p \right\}$

Proof:  $(\subseteq)$   $v \in T_p^{\text{sub}} M \Rightarrow v = J_\psi(\tilde{\varphi}^{-1}(p)) x$  for  $x \in \mathbb{R}^k$ ,  $\psi$  local parameterisation  
 $\Rightarrow v = J_\psi(\tilde{\gamma}(0)) \tilde{\gamma}'(0)$  with  $\tilde{\gamma}(t) = \tilde{p} + tx$ ,  $\tilde{\gamma}: (-\epsilon, \epsilon) \rightarrow \mathbb{R}^k$   
 $= \frac{d}{dt} (\underbrace{\psi \circ \tilde{\gamma}}_\gamma) \Big|_{t=0} = \gamma'(0)$

$(\supseteq)$  Take:  $\gamma: (-\epsilon, \epsilon) \rightarrow M$  differentiable with  $\gamma(0) = p$




$$\gamma'(0) = \frac{d}{dt} (\psi \circ \tilde{\gamma}) \Big|_{t=0} = J_\psi(\tilde{\gamma}(0)) \tilde{\gamma}'(0) = J_\psi(\tilde{\varphi}^{-1}(p)) x \in T_p^{\text{sub}} M$$

# Manifolds - Part 21

$$T_p^{\text{sub}} M \rightsquigarrow T_p M$$

for  $M \subseteq \mathbb{R}^n$  smooth submanifold      for  $M$  smooth manifold



Definition:  $C_p(M) := \{ \gamma : (-\epsilon, \epsilon) \rightarrow M \mid \gamma \text{ differentiable with } \gamma(0) = p \}$

$$\gamma \sim \alpha \iff (h \circ \gamma)'(0) = (h \circ \alpha)'(0)$$

for a chart  $(U, h)$ .

equivalent class:  $[\gamma]_{\sim} := \{ \alpha \mid \gamma \sim \alpha \}$  represents **tangent vector**

$$T_p M := C_p(M) / \sim \quad (\text{set of all equivalence classes})$$

tangent space of the manifold  $M$

Result:

- For a submanifold  $T_p^{\text{sub}} M \xleftrightarrow{\text{bijection}} T_p M$
- $\gamma'(0) \xleftrightarrow{\text{bijection}} [\gamma]_{\sim}$

- $T_p M$  is a vector space with the operations:

$$v + w := h_*^{-1} (h_*(v) + h_*(w)) \quad \text{with } h_*: [\gamma]_{\sim} \mapsto (h \circ \gamma)'(0) \in \mathbb{R}^k$$

$$\lambda \cdot v := h_*^{-1} (\lambda \cdot h_*(v))$$

# Manifolds - Part 22

smooth manifold  $M$  of dimension  $n$ ,  $p \in M$ .





chart  $(U, h)$ :




defined by:  
 $h_* : T_p M \rightarrow \mathbb{R}^n$   
 $[\gamma] \mapsto (h \circ \gamma)'(0)$   
 linear + bijective  
 $\psi_* := h_*^{-1}$

Definition: coordinate basis (standard basis with respect to  $(U, h)$ ):

For  $(U, h)$  and  $p \in U$ , we define:  $\partial_j := \psi_*(e_j)$

where  $(e_1, e_2, \dots, e_n)$  is the standard basis of  $\mathbb{R}^n$

Remember: For submanifolds:




$(\partial_1, \partial_2, \dots, \partial_n)$  is essentially  $\left( \frac{\partial \psi}{\partial x_1}(\tilde{p}), \frac{\partial \psi}{\partial x_2}(\tilde{p}), \dots, \frac{\partial \psi}{\partial x_n}(\tilde{p}) \right)$


Soon:  $f: M \rightarrow N$  smooth  $\rightsquigarrow$   $df_p: T_p M \rightarrow T_p N$  differential



# Manifolds - Part 23



Definition: tangent bundle  $TM := \bigsqcup_{p \in M} T_p M := \bigcup_{p \in M} \{p\} \times T_p M$   
 ↪ smooth manifold of dimension  $2 \cdot \dim(M)$




Definition: differential of f at point p

$$df_p : T_p M \longrightarrow T_{f(p)} N$$


$$[\gamma] \longmapsto [f \circ \gamma]$$

differential:  $df : p \longmapsto df_p$

Example for submanifolds  $M, N \subseteq \mathbb{R}^n$  smooth submanifolds



$f$   
smooth



bijection  
 $T_{f(p)} N \cong T_{f(p)}^{sub} N$

$$[\gamma] \xrightarrow{df_p} [f \circ \gamma] \stackrel{\text{bijection}}{=} (f \circ \gamma)'(0)$$


Example:  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  (smooth map)

$$df_p([\gamma]) \stackrel{\text{bijection}}{=} (f \circ \gamma)'(0) = J_f(\underbrace{\gamma(0)}_p) \underbrace{\gamma'(0)}_{\text{tangent vector}}$$

= directional derivative of  $f$  along  $[\gamma]$  at  $p$

# Manifolds - Part 24

Differential in local charts?



Choose:  $[\gamma] \in T_p M$  :

$$\begin{aligned}
 dk_{f(p)}(df_p([\gamma])) &= dk_{f(p)}([f \circ \gamma]) \\
 &= [k \circ f \circ \gamma] \stackrel{\text{bijection}}{=} (k \circ f \circ \gamma)'(0) \\
 &= (\tilde{f} \circ h \circ \gamma)'(0) \\
 &\stackrel{\text{ordinary chain rule}}{=} J_{\tilde{f}}(h(p)) (h \circ \gamma)'(0) \\
 &\stackrel{\text{bijection}}{=} J_{\tilde{f}}(h(p)) [h \circ \gamma] \\
 &= J_{\tilde{f}}(h(p)) dh_p([\gamma])
 \end{aligned}$$

Remember:

$$\begin{aligned}
 f &= k^{-1} \circ \tilde{f} \circ h \\
 df &= dk^{-1} J_{\tilde{f}} dh
 \end{aligned}$$

# Manifolds - Part 25

Recall:  $p \in M$ ,  $(U, h)$ : coordinate basis  $(\partial_1, \dots, \partial_n)$  of  $T_p M$   
 $\varphi = h^{-1}$ ,  $\partial_j := \varphi_*(e_j) = d\varphi_{h(p)}(e_j)$

defined by:  
 $h_*: T_p M \rightarrow \mathbb{R}^n$   
 $[\gamma] \mapsto (h \circ \gamma)'(0)$   
 linear + bijective  
 $\varphi_* := h_*^{-1}$

Directional derivative:  $f: M \rightarrow \mathbb{R}$  smooth

$$(\partial_j f)(p) := df_p(\partial_j)$$

$$= df_p(d\varphi_{h(p)}(e_j))$$


$$= [f \circ \varphi \circ \tilde{\gamma}]$$

bijection


$$= (f \circ \varphi \circ \tilde{\gamma})'(0)$$

chain rule

$$= J_{f \circ \varphi}(h(p)) \underbrace{\tilde{\gamma}'(0)}_{e_j} = \frac{\partial (f \circ \varphi)}{\partial x_j}(h(p))$$



Example:



$$\partial_1 = d\varphi_{h(z)}(e_1) = [\varphi \circ \tilde{\gamma}]', \quad \tilde{\gamma}(t) = h(z) + t$$

$$= (\varphi \circ \tilde{\gamma})'(0) = \frac{d}{dt} \Big|_{t=0} e^{i(s+t)} = i \cdot e^{is}$$

$$\tilde{\partial}_1 = d\psi_{k(f(z))}(e_1)$$

$$= (\psi \circ \tilde{\gamma})'(0) \quad \tilde{\gamma}(t) = k(z^2) + t$$

$$= i \cdot e^{2is} \quad \downarrow$$


$$(e^{is})^2$$

map  $\tilde{f}$ :  $s \xrightarrow{\varphi} e^{is} \xrightarrow{f} (e^{is})^2 \xrightarrow{k} 2s$

$$J_{\tilde{f}}(s) = 2$$

differential of f:  $df_z(\partial_1) \stackrel{\text{last video}}{=} dk_{z^2}^{-1} \underbrace{J_{\tilde{f}}(h(p))}_2 \underbrace{dh_z(\partial_1)}_{e_1} = 2 \cdot dk_{z^2}^{-1}(e_1) = 2 \cdot \tilde{\partial}_1$

# Manifolds - Part 26



## Introduction to Ricci calculus / tensor calculus

↳ calculating in coordinates

↳ positions of indices matter  
(superscripts, subscripts)

our language	Ricci calculus
components of a given chart $(U, h)$ , $h: U \rightarrow \mathbb{R}^n$	$h^j: U \rightarrow \mathbb{R}$ coordinates or simply: $x^1, x^2, \dots, x^n$
coordinate basis of $T_p M$ : $\partial_j := \psi_*(e_j)$	$\frac{\partial}{\partial x^1}, \frac{\partial}{\partial x^2}, \dots, \frac{\partial}{\partial x^n}$
tangent vector $[v] \in T_p M$ : $v_1 \partial_1 + v_2 \partial_2 + \dots + v_n \partial_n$	$v^1 \frac{\partial}{\partial x^1} + \dots + v^n \frac{\partial}{\partial x^n} =: v^j \frac{\partial}{\partial x^j}$ (Einstein summation convention) <u>contravariant</u> vector
inner product on $T_p M$ : $\langle v, w \rangle \in \mathbb{R}$	$v^j \underbrace{g_{jk}} w^k$ → tensor

Later:

dual to a contravariant vector:

$$v_j \underbrace{dx^j}$$


↳ one-form ( $\rightsquigarrow$  linear map)

$$dx_j(\partial_k) = \begin{cases} 1, & j = k \\ 0, & j \neq k \end{cases}$$

$$= \delta_{jk}$$


Kronecker delta

$$dx^j\left(\frac{\partial}{\partial x^k}\right) = \delta^j_k$$



## Manifolds - Part 27

Recall:



$T_p M$   $n$ -dimensional vector space

$$\text{Define: } T_p^* M := (T_p M)^*$$

$$= \{ \alpha: T_p M \rightarrow \mathbb{R} \text{ linear} \}$$

$$\leadsto dx_{j,p}: T_p M \rightarrow \mathbb{R}$$

$$dx_{j,p}(\partial_k) = \delta_{jk} \quad \text{linear map!}$$

differential form: map  $\omega$  defined on  $M$  such that  $\omega(p) \in T_p^* M$   
(one-form)

$$dx_j: p \mapsto dx_{j,p} \in T_p^* M$$

Some multilinear algebra:  $\text{Alt}^k(V) := \left\{ \alpha: \underbrace{V \times \dots \times V}_{k\text{-times}} \rightarrow \mathbb{R} \text{ multilinear (k-linear)} \right\}$

+ alternating

$$\alpha(v_1, \dots, v_k) = 0$$

if  $(v_1, \dots, v_k)$

linearly dependent


Example:  $\alpha \in \text{Alt}^2(V)$ ,  $\alpha(v_1, v_2) = -\alpha(v_2, v_1)$

$$\det \in \text{Alt}^2(\mathbb{R}^2)$$

$\alpha \in \text{Alt}^k(V)$  is called an alternating  $k$ -form on  $V$

Remember:  $\text{Alt}^1(V) = V^*$  (dual space of  $V$ )

$$\text{Alt}^0(V) = \mathbb{R}$$



## Manifolds - Part 28

Wedge product:  $\wedge$  multiplication defined for  $\alpha \in \text{Alt}^k(V)$ ,  $\beta \in \text{Alt}^s(V)$

$$\begin{aligned}\wedge : \text{Alt}^k(V) \times \text{Alt}^s(V) &\longrightarrow \text{Alt}^{k+s}(V) \\ (\alpha, \beta) &\longmapsto \alpha \wedge \beta\end{aligned}$$

$$\overset{(k+s)\text{-linear}}{\curvearrowright} (\alpha \wedge \beta)(v_1, \dots, v_{k+s}) \neq \alpha(v_1, \dots, v_k) \cdot \beta(v_{k+1}, \dots, v_{k+s})$$

not a possible definition!  
(not alternating)

Definition: For  $\alpha \in \text{Alt}^k(V)$ ,  $\beta \in \text{Alt}^s(V)$ , we define  $\alpha \wedge \beta \in \text{Alt}^{k+s}(V)$  by:

$$(\alpha \wedge \beta)(v_1, \dots, v_{k+s}) := \frac{1}{k! \cdot s!} \sum_{\sigma \in S_{k+s}} \text{sgn}(\sigma) \alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \beta(v_{\sigma(k+1)}, \dots, v_{\sigma(k+s)})$$

Examples: (a)  $\alpha, \beta \in \text{Alt}^1(V) = V^*$ :

$$(\alpha \wedge \beta)(u, v) = \alpha(u)\beta(v) - \alpha(v)\beta(u)$$

$$(b) \alpha, \beta \in \text{Alt}^1(\mathbb{R}^3), \quad \alpha\left(\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}\right) = x_1, \quad \beta\left(\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}\right) = x_2 = \underbrace{(0, 1, 0)}_{\text{identified with } \beta} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

$$(\alpha \wedge \beta)\left(\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}\right) = x_1 y_2 - y_1 x_2 = \left\langle \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \underbrace{\begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}}_{\text{identified with } \alpha \wedge \beta} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \right\rangle$$

Properties:

$$(a) \quad \alpha \wedge \beta = (-1)^{k \cdot s} \beta \wedge \alpha \quad (\text{anticommutative})$$

$$(b) \quad (\alpha + \alpha') \wedge \beta = \alpha \wedge \beta + \alpha' \wedge \beta$$

$$(\lambda \alpha) \wedge \beta = \lambda (\alpha \wedge \beta) \quad (\text{bilinear})$$

$$(c) \quad \alpha \wedge (\beta \wedge \gamma) = (\alpha \wedge \beta) \wedge \gamma \quad (\text{associative})$$


(d) For a linear map  $f: W \rightarrow V$  and  $\alpha \in \text{Alt}^k(V)$  define:

$$\text{pullback} \quad (f^* \alpha)(w_1, \dots, w_k) := \alpha(f(w_1), \dots, f(w_k))$$

("natural")

$$f^*(\alpha \wedge \beta) = f^* \alpha \wedge f^* \beta$$





## Manifolds - Part 29

$M$  smooth manifold of dimension  $n \Rightarrow T_p M$   $n$ -dimensional vector space

Definition:

$$\omega : M \longrightarrow \bigcup_{p \in M} \text{Alt}^k(T_p M)$$

$$p \longmapsto \omega_p = \omega(p) \in \text{Alt}^k(T_p M)$$


is called a  $k$ -form on  $M$ .

We also define:  $\omega \wedge \eta$  as  $(\omega \wedge \eta)(p) := \omega(p) \wedge \eta(p)$

$$f^* \omega \quad \text{as} \quad (f^* \omega)(p) := (df_p)^* \omega(p)$$

$$f : N \longrightarrow M \text{ smooth}$$

Basis elements:



basis of  $T_p M$  :  $(\partial_1, \partial_2, \dots, \partial_n)$  with  $\partial_j := \varphi_*(e_j) = d\varphi_{h(p)}(e_j)$

basis of  $(T_p M)^* = \text{Alt}^1(T_p M)$  :  $(dx_p^1, dx_p^2, \dots, dx_p^n)$

$$\text{defined by: } dx_p^j(\partial_k) = \delta_k^j = \begin{cases} 1 & , j=k \\ 0 & , j \neq k \end{cases}$$

Proposition: A basis of  $\text{Alt}^k(T_p M)$  is given by:

$$(dx_p^{\mu_1} \wedge dx_p^{\mu_2} \wedge \dots \wedge dx_p^{\mu_k})_{\mu_1 < \mu_2 < \dots < \mu_k}$$

Example:  $\dim(M) = 3$ ,  $\text{Alt}^2(T_p M)$ :

$$(dx_p^1 \wedge dx_p^2, dx_p^1 \wedge dx_p^3, dx_p^2 \wedge dx_p^3)$$

Conclusion: Each  $k$ -form on  $M$  can locally be written as:

$$\omega(p) = \sum_{\mu_1 < \dots < \mu_k} \omega_{\mu_1, \mu_2, \dots, \mu_k}(p) \cdot dx_p^{\mu_1} \wedge dx_p^{\mu_2} \wedge \dots \wedge dx_p^{\mu_k}$$

$$\omega_{\mu_1, \mu_2, \dots, \mu_k} : U \longrightarrow \mathbb{R} \quad \text{component functions}$$


Definition: • If all component functions are differentiable at  $p$ ,  
then  $\omega$  is differentiable at  $p$ .

• If  $\omega$  is differentiable at all  $p \in M$ ,

then  $\omega$  is called a differential form on  $M$ .

$$\omega \in \Omega^k(M)$$

$$\Omega^0(M) := C^\infty(M)$$



## Manifolds - Part 30



differential form on a manifold:  $\omega \in \Omega^k(M)$   $\leftarrow$   $k$ -form on  $M$   
+ differentiable

$$\omega(p) = \sum_{\mu_1 < \dots < \mu_k} \omega_{\mu_1, \mu_2, \dots, \mu_k}(p) \cdot dx_p^{\mu_1} \wedge dx_p^{\mu_2} \wedge \dots \wedge dx_p^{\mu_k}$$

Examples:

(a)

$$M = \mathbb{R}^2$$



$$\partial_k = e_k$$

$$dx_p^j(\partial_k) = \delta^j_k$$

identify:  $\partial_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $dx_p^1 = (1, 0)$

$$\partial_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad dx_p^2 = (0, 1)$$


$$\begin{aligned} (dx_p^1 \wedge dx_p^2) \left( \begin{matrix} a_1 & a_2 \\ \parallel & \parallel \\ \begin{pmatrix} a_{1,1} \\ a_{2,1} \end{pmatrix} & \begin{pmatrix} a_{1,2} \\ a_{2,2} \end{pmatrix} \end{matrix} \right) &= \sum_{\sigma \in S_2} \text{sgn}(\sigma) dx_p^1(a_{\sigma(1)}) dx_p^2(a_{\sigma(2)}) \\ &= \sum_{\sigma \in S_2} \text{sgn}(\sigma) a_{1,\sigma(1)} a_{2,\sigma(2)} = \det \begin{pmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{pmatrix} \end{aligned}$$

(b) Each  $\omega \in \Omega^n(\mathbb{R}^n)$  can be written as:

$$\omega(p) = \omega_{1,2,\dots,n}(p) dx_p^1 \wedge dx_p^2 \wedge \dots \wedge dx_p^n$$

$$= \omega_{1,2,\dots,n}(p) \det \begin{pmatrix} | & | & \dots & | \\ | & | & \dots & | \\ | & | & \dots & | \end{pmatrix}$$

(c)  $M = \mathbb{R}^2$



$\varphi$  given by polar coordinates  $\varphi(r, \theta) = \begin{pmatrix} r \cdot \cos(\theta) \\ r \cdot \sin(\theta) \end{pmatrix}$

$(r, \theta)$

$$\partial_j := \varphi_*(e_j) = J_\varphi(\tilde{p})(e_j)$$

$$\partial_1(r, \theta) = \frac{\partial \varphi}{\partial r}(r, \theta) = \begin{pmatrix} \cos(\theta) \\ \sin(\theta) \end{pmatrix}$$


$$\partial_2(r, \theta) = \frac{\partial \varphi}{\partial \theta}(r, \theta) = \begin{pmatrix} -r \cdot \sin(\theta) \\ r \cdot \cos(\theta) \end{pmatrix}$$

corresponding 1-forms:  $d\Gamma_p = (\cos(\theta), \sin(\theta)) = \frac{1}{\sqrt{x^2+y^2}}(x, y)$

for  $p = (x, y)$   $d\theta_p = \frac{1}{r}(-\sin(\theta), \cos(\theta)) = \frac{1}{x^2+y^2}(-y, x)$

2-form:  $(d\Gamma_p \wedge d\theta_p)(e_1, e_2) = d\Gamma_p(e_1)d\theta_p(e_2) - d\Gamma_p(e_2)d\theta_p(e_1)$   
 $= \frac{1}{r}(\cos(\theta))^2 - \frac{1}{r} \cdot (-1)(\sin(\theta))^2$   
 $= \frac{1}{r}$

$$\Rightarrow r d\Gamma_p \wedge d\theta_p = \det \begin{pmatrix} | & | \\ | & | \end{pmatrix} = dx_p \wedge dy_p$$



# Manifolds - Part 31

vector space  $\leftarrow$  orientation

for example:  $\mathbb{R}^n$  with basis:  $\mathcal{B} = (e_1, e_2, \dots, e_n)$

change-of-basis matrix  $T_{\mathcal{C} \leftarrow \mathcal{B}}$



two cases:

$\det(T_{\mathcal{C} \leftarrow \mathcal{B}}) > 0$  : positively orientated

$\det(T_{\mathcal{C} \leftarrow \mathcal{B}}) < 0$  : negatively orientated

$\mathcal{C} = (c_1, c_2, \dots, c_n)$


$\Rightarrow$  two equivalence classes for bases



Remember:  $V$  finite-dimensional vector space + one chosen equivalence class

$\rightsquigarrow$  orientation  $(V, \text{or})$

Orientations for manifolds:




Definition: A smooth manifold  $M$  is called orientable if there is a family of orientations for the tangent spaces  $\{(T_p M, or_p)\}$  such that


$$\forall p \in M \exists (U, h) \forall x \in U: (\partial_1^{(h)}(x), \partial_2^{(h)}(x), \dots, \partial_n^{(h)}(x)) \in or_x$$


Example: (a) If  $M$  has an atlas with one chart  $(M, h)$ , then  $M$  is orientable.

(b) Möbius strip:




after running  
around the strip:





## Manifolds - Part 32



orientable manifold  $M$

Fact: Let  $M$  be an  $n$ -dim smooth manifold. Then the following claims are equivalent:

(a)  $M$  is orientable: We have  $\{(T_p M, or_p)\}$  such that


$$\forall p \in M \exists (U, h) \forall x \in U: (\partial_1^{(h)}(x), \partial_2^{(h)}(x), \dots, \partial_n^{(h)}(x)) \in or_x$$

(b) There is an atlas for  $M$  collection of charts that cover the manifold

such that all transition maps

$w: \text{green shape} \rightarrow \text{green shape}$  satisfy:


$$\det(J_w(x)) > 0$$



(c) There is a differential form (volume form)

$$\omega \in \Omega^n(M) \quad \text{with} \quad \omega(p) \neq 0 \quad \text{for all } p \in M.$$

Proof: (a)  $\Leftrightarrow$  (b)



We have:  $\psi_* ( \underbrace{J_\omega(\tilde{p}) e_1}_{\text{first column of Jacobian}} ) = \psi_* (e_1)$

$$= \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{pmatrix} = \sum_j \lambda_j e_j$$

$$\Rightarrow \sum_{j=1}^n \lambda_j \underbrace{\psi_*(e_j)}_{\partial_j^{(k)}(p)} = \underbrace{\psi_*(e_1)}_{\partial_1^{(h)}(p)} \quad (*)$$

Change-of-basis matrix:  $\mathcal{B} = (\partial_1^{(h)}(p), \dots, \partial_n^{(h)}(p)) \xrightarrow{T_{\mathcal{C} \leftarrow \mathcal{B}}} \mathcal{C} = (\partial_1^{(k)}(p), \dots, \partial_n^{(k)}(p))$


$$\Rightarrow (*) \quad T_{\mathcal{C} \leftarrow \mathcal{B}} = \begin{pmatrix} \lambda_1 & & \\ & \lambda_2 & \dots \\ & \vdots & \\ & & \lambda_n \end{pmatrix} = J_\omega(\tilde{p})$$

Hence:


$$\det(T_{\mathcal{C} \leftarrow \mathcal{B}}) > 0 \quad \Leftrightarrow \quad \det(J_\omega(x)) > 0$$

$$(a) \quad \Leftrightarrow \quad (b)$$






# Manifolds - Part 33



In  $\mathbb{R}^n$ : inner product  $\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$

write:  $g(x, y) = \langle x, y \rangle$




Definition:  $M$  smooth manifold. If we have an inner product  $g_p$  on  $T_p M$  for all  $p \in M$  and  $p \mapsto g_p$  smooth, then:

$g: p \mapsto g_p$  is called a Riemannian metric and

$(M, g)$  is called a Riemannian manifold.

What does smooth mean?



$$g_x(\partial_i^{(h)}(x), \partial_j^{(h)}(x)) =: g_{ij}^{(h)}(x) \quad \left( \partial_1^{(h)}(x), \partial_2^{(h)}(x), \dots, \partial_n^{(h)}(x) \right)$$


maps:  $U \rightarrow \mathbb{R}^n$  smooth!

$$x \mapsto g_{ij}^{(h)}(x) \quad \text{for all } i, j, (U, h)$$

(Einstein summation convention)

In local coordinates:  $g_x(\cdot, \cdot) = g_{ij}^{(h)}(x) dx_x^i(\cdot) dx_x^j(\cdot)$

Hence:  $g_x$  can be seen as a symmetric matrix:  $G = \left( g_{ij}^{(h)}(x) \right)_{ij}$




## Manifolds - Part 34

Riemannian metric:

$$g: P \mapsto g_P \leftarrow \text{inner product on } T_P M$$

smooth


Submanifolds in  $\mathbb{R}^N$ :



Note:  $T_P M \cong T_P^{\text{sub}} M = \text{span}\left(\frac{\partial \varphi}{\partial x_1}, \dots, \frac{\partial \varphi}{\partial x_n}\right)$

$$g_{ij}^{(h)}(p) = \left\langle \frac{\partial \varphi}{\partial x_i}(\tilde{p}), \frac{\partial \varphi}{\partial x_j}(\tilde{p}) \right\rangle_{\text{standard}}$$

Examples: (a) 1-dimensional submanifold in  $\mathbb{R}^N$



$$g_{11}^{(h)}(p) = \left\langle \varphi'(t), \varphi'(t) \right\rangle_{\text{standard}} = \|\varphi'(t)\|_{\text{standard}}^2$$

length:  $\int_a^b \|\varphi'(t)\|_{\text{standard}} dt = \int_a^b \sqrt{\det(G)} dt$

(b)  $S^2 \subseteq \mathbb{R}^3$  has parameterization given by spherical coordinates:


$$\Phi(\theta, \varphi) = \begin{pmatrix} \sin(\theta) \cos(\varphi) \\ \sin(\theta) \sin(\varphi) \\ \cos(\theta) \end{pmatrix}$$

$$\Rightarrow \text{two tangent vectors: } \frac{\partial \Phi}{\partial \theta} = \begin{pmatrix} \cos(\theta) \cos(\varphi) \\ \cos(\theta) \sin(\varphi) \\ -\sin(\theta) \end{pmatrix}$$

$$\frac{\partial \Phi}{\partial \varphi} = \begin{pmatrix} -\sin(\theta) \sin(\varphi) \\ \sin(\theta) \cos(\varphi) \\ 0 \end{pmatrix}$$

$$\Rightarrow G = \begin{pmatrix} 1 & 0 \\ 0 & \sin^2(\theta) \end{pmatrix} \rightsquigarrow \sqrt{\det(G)} = |\sin(\theta)|$$

$$\text{volume form: } \sqrt{\det(G)} \, d\theta \wedge d\varphi$$



## Manifolds - Part 35

We already know: An orientable  $n$ -dimensional manifold  $M$  has a non-trivial volume form  $\omega \in \Omega^n(M)$ .



Definition:  $M$  orientable Riemannian manifold of dimension  $n$ .

Then the canonical volume form  $\omega_M \in \Omega^n(M)$  is defined by:

If  $(v_1, v_2, \dots, v_n)$  is a positively orientated basis of  $T_p M$  and an orthonormal basis of  $T_p M$  (ONB),  $g_p(v_i, v_j) = \delta_{ij}$

then:  $\omega_M(p)(v_1, v_2, \dots, v_n) = 1$

Proposition:  $(M, g)$  orientable Riemannian manifold of dimension  $n$ .



Let  $(U, h)$  be a chart such that the basis

$$(\partial_1^{(h)}(x), \partial_2^{(h)}(x), \dots, \partial_n^{(h)}(x))$$

is positively orientated for all  $x \in U$ .

$$\omega_M(x) = \sqrt{\det(G)} dx_x^1 \wedge dx_x^2 \wedge \dots \wedge dx_x^n$$

where  $G_{ij} := g_x(\partial_i^{(h)}(x), \partial_j^{(h)}(x))$

determinant of Gram/ Gramian


dual basis

Proof:

$$\begin{array}{ccc}
 (\partial_1^{(h)}(x), \partial_2^{(h)}(x), \dots, \partial_n^{(h)}(x)) & \xrightarrow{\text{Gram-Schmidt}} & (v_1, v_2, \dots, v_n) \text{ ONB} \\
 \uparrow \text{positively orientated} & \xleftarrow{f} & \uparrow \text{positively orientated} \\
 & \text{linear map} & 
 \end{array}$$


$$\begin{aligned}
 \text{Then: } \omega_M(x) (\partial_1^{(h)}(x), \partial_2^{(h)}(x), \dots, \partial_n^{(h)}(x)) & \\
 = \omega_M(x) (f(v_1), f(v_2), \dots, f(v_n)) &= f^* \omega_M(x) (v_1, \dots, v_n) \\
 = \det(f) \underbrace{\omega_M(x) (v_1, \dots, v_n)}_{=1} & \\
 g_x(\partial_i^{(h)}(x), \partial_j^{(h)}(x)) = g_x(f(v_i), f(v_j)) &
 \end{aligned}$$

$$\begin{array}{ccc}
 \begin{array}{ccc}
 \boxed{\begin{array}{c} \uparrow \\ \rightarrow \\ v_i \end{array}} & \xrightarrow{f} & \boxed{\begin{array}{c} \uparrow \\ \rightarrow \\ v_j \end{array}} \\
 \downarrow \Phi & & \downarrow \Phi \\
 \boxed{\begin{array}{c} \uparrow \\ \rightarrow \\ e_i \end{array}} & \xrightarrow{A} & \boxed{\begin{array}{c} \uparrow \\ \rightarrow \\ e_j \end{array}}
 \end{array} & & 
 \begin{aligned}
 &= g_x(\Phi^{-1} A \Phi(v_i), \Phi^{-1} A \Phi(v_j)) \\
 &= \langle \underbrace{A \Phi(v_i)}_{e_i}, \underbrace{A \Phi(v_j)}_{e_j} \rangle_{\text{standard}} = (A^T A)_{ij} \\
 \Rightarrow \det(G) &= \det(A)^2 \quad \square
 \end{aligned}
 \end{array}$$




## Manifolds - Part 36

$M$  orientable Riemannian manifold of dimension  $n$ .

↳ canonical volume form  $\omega_M(x) = \sqrt{\det(G)} dx_x^1 \wedge \dots \wedge dx_x^n$    $\square \subseteq \mathbb{R}^n$

Examples: (a)  $S^2 \subseteq \mathbb{R}^3$  has parameterization given by spherical coordinates:

$$\Phi(\theta, \varphi) = \begin{pmatrix} \sin(\theta) \cos(\varphi) \\ \sin(\theta) \sin(\varphi) \\ \cos(\theta) \end{pmatrix}$$




$$\Rightarrow G = \begin{pmatrix} 1 & 0 \\ 0 & \sin^2(\theta) \end{pmatrix}$$

$$\Rightarrow \omega_M(x) = \sin(\theta) d\theta \wedge d\varphi$$

(b) Graph surface:  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$   $C^\infty$ -function

$$M := \{(x, f(x)) \mid x \in \mathbb{R}^2\}$$

2-dim. submanifold in  $\mathbb{R}^3$



Use parameterization:  $\varphi: x \mapsto (x, f(x))$ ,  $h: (x, f(x)) \mapsto x$

$$\text{tangent vectors: } \partial_1^{(h)}(p) \stackrel{\text{identify}}{=} \frac{\partial \varphi}{\partial x_1}(x) = \begin{pmatrix} 1 \\ 0 \\ \frac{\partial f}{\partial x_1}(x) \end{pmatrix}$$

$$\partial_2^{(h)}(p) \stackrel{\text{identify}}{=} \frac{\partial \varphi}{\partial x_2}(x) = \begin{pmatrix} 0 \\ 1 \\ \frac{\partial f}{\partial x_2}(x) \end{pmatrix}$$

$$g_{ij}^{(h)}(p) = \left\langle \frac{\partial \varphi}{\partial x_i}(x), \frac{\partial \varphi}{\partial x_j}(x) \right\rangle_{\text{standard}} = \begin{cases} \frac{\partial f}{\partial x_i} \cdot \frac{\partial f}{\partial x_j}, & i \neq j \\ 1 + \frac{\partial f}{\partial x_i} \cdot \frac{\partial f}{\partial x_j}, & i = j \end{cases}$$


$$\Rightarrow G = \begin{pmatrix} 1 + \left(\frac{\partial f}{\partial x_1}\right)^2 & \frac{\partial f}{\partial x_1} \cdot \frac{\partial f}{\partial x_2} \\ \frac{\partial f}{\partial x_1} \cdot \frac{\partial f}{\partial x_2} & 1 + \left(\frac{\partial f}{\partial x_2}\right)^2 \end{pmatrix}$$

$$\det(G) = 1 + \left(\frac{\partial f}{\partial x_1}\right)^2 + \left(\frac{\partial f}{\partial x_2}\right)^2$$


Canonical volume form:  $\omega_M(p) = \sqrt{1 + \left(\frac{\partial f}{\partial x_1}\right)^2 + \left(\frac{\partial f}{\partial x_2}\right)^2} dx_1^1 \wedge dx_1^2$

Interesting fact:  $\left\| \partial_1^{(h)}(p) \times \partial_2^{(h)}(p) \right\|_{\text{standard}} = \left\| \begin{pmatrix} 1 \\ 0 \\ \frac{\partial f}{\partial x_1}(x) \end{pmatrix} \times \begin{pmatrix} 0 \\ 1 \\ \frac{\partial f}{\partial x_2}(x) \end{pmatrix} \right\|_{\text{standard}}$

$$= \left\| \begin{pmatrix} -\frac{\partial f}{\partial x_1} \\ -\frac{\partial f}{\partial x_2} \\ 1 \end{pmatrix} \right\|_{\text{standard}} = \sqrt{\det(G)}$$



## Manifolds - Part 37



$M \subseteq \mathbb{R}^3$  orientable Riemannian manifold of dimension 2

length of  $N \iff$  canonical volume form

Definition: Let  $\tilde{M}$  be a Riemannian manifold and  $M \subseteq \tilde{M}$ .


A map  $N: M \rightarrow T\tilde{M}$

$$p \mapsto N(p) \in T_p \tilde{M}$$

$$\text{and } N(p) \in (T_p M)^\perp \setminus \{0\}$$

is called a normal vector field.


(see  $T_p M \subseteq T_p \tilde{M}$ )  
(orthogonal w.r.t.  $g_p$ )



We call it continuous at  $p$  if for a chart  $(U, h)$  of  $\tilde{M}$  with  $p \in U$  holds:

$$N(x) = \sum_i a_i(x) \cdot \partial_i^{(h)}(x)$$

continuous functions  $U \rightarrow \mathbb{R}$




We call it a continuous unit normal vector field if

- $N$  is continuous at every  $p \in M$
- $\|N(x)\| = \sqrt{g_x(N(x), N(x))} = 1$  for all  $x \in M$ .

Important fact:

$M \subseteq \mathbb{R}^n$   $(n-1)$ -dimensional submanifold:

(a)  $M$  is orientable  $\iff M$  has a continuous unit normal vector field



continuous normal vector field not possible




(b) If  $N$  is a continuous unit normal vector field, then:

canonical volume form  $\rightarrow \omega_M = N \lrcorner \det$

means:


$$\omega_M(x)(v_1, \dots, v_{n-1}) = \det(N(x), v_1, \dots, v_{n-1})$$



Example:

$$S^2 \subseteq \mathbb{R}^3,$$

$$N(x) = x$$



parameterization:

$$\Phi(\theta, \psi) = \begin{pmatrix} \sin(\theta) \cos(\psi) \\ \sin(\theta) \sin(\psi) \\ \cos(\theta) \end{pmatrix}$$

$$\sqrt{\det(G)} = \omega_M(x)(\partial_1^{(h)}(x), \partial_2^{(h)}(x)) = \det(N(x), \partial_1^{(h)}(x), \partial_2^{(h)}(x))$$

$$= \det \begin{pmatrix} \sin(\theta) \cos(\psi) & \cos(\theta) \cos(\psi) & -\sin(\theta) \sin(\psi) \\ \sin(\theta) \sin(\psi) & \cos(\theta) \sin(\psi) & \sin(\theta) \cos(\psi) \\ \cos(\theta) & -\sin(\theta) & 0 \end{pmatrix}$$

$$= \sin(\theta)$$