



Measure Theory - Part 14

Radon-Nikodym theorem and Lebesgue's decomposition theorem

$(X, \mathcal{A}, \lambda)$ measure space
 $\stackrel{\parallel}{=} \mathbb{R} \quad \stackrel{\parallel}{=} \mathcal{B}(\mathbb{R}) \quad \stackrel{\parallel}{=} \lambda$
 (Lebesgue measure) reference measure $(\lambda([a, b]) = b - a)$

Another measure $\mu: \mathcal{B}(\mathbb{R}) \rightarrow [0, \infty]$

Definitions: (a) μ is called absolutely continuous (w.r.t λ)

if $\lambda(A) = 0 \Rightarrow \mu(A) = 0$
 for all $A \in \mathcal{B}(\mathbb{R})$. One writes: $\mu \ll \lambda$

(b) μ is called singular (w.r.t. λ)

if there is $N \in \mathcal{B}(\mathbb{R})$ with $\lambda(N) = 0$ and $\mu(\mathbb{R} \setminus N) = 0$.

One writes: $\mu \perp \lambda$

Example: δ_0 Dirac measure ($\delta_0(\{0\}) = 1$)
 $\Rightarrow \delta_0 \perp \lambda$ (Choose $N = \{0\}$)

Theorem: $\mu: \mathcal{B}(\mathbb{R}) \rightarrow [0, \infty]$ (σ -finite)

(a) There are two measures (uniquely determined)

Lebesgue's
decomposition
theorem

$$\mu_{ac}, \mu_s: \mathcal{B}(\mathbb{R}) \rightarrow [0, \infty]$$

with $\mu = \mu_{ac} + \mu_s$, $\mu_{ac} \ll \lambda$, $\mu_s \perp \lambda$.

(b) There is a measurable map $h: \mathbb{R} \rightarrow [0, \infty)$

Radon-
Nikodym
theorem

with $\mu_{ac}(A) = \int_A h d\lambda$
 for all $A \in \mathcal{B}(\mathbb{R})$. density