



The Bright Side of Mathematics

The following pages cover the whole Multidimensional Integration course of the Bright Side of Mathematics. Please note that the creator lives from generous supporters and would be very happy about a donation. See more here: <https://tbsom.de/support>

Have fun learning mathematics!




Multidimensional Integration - Part 1



Lebesgue measure on \mathbb{R} :

$$\text{length: } \mu([a,b]) = b - a \quad a \leq b$$

Carathéodory's Extension Theorem




$$\psi: \mathcal{P}(\mathbb{R}) \rightarrow [0, \infty] \quad \text{outer measure}$$

$$\mathcal{L}(\mathbb{R}) = \mathcal{A}_\psi \quad \sigma\text{-algebra of Lebesgue-measurable sets}$$

$$\lambda: \mathcal{L}(\mathbb{R}) \rightarrow [0, \infty] \quad \text{Lebesgue measure on } \mathbb{R}$$

measure


Properties:


- $\lambda(\emptyset) = 0$
- $\lambda\left(\bigcup_{j=1}^{\infty} A_j\right) = \sum_{j=1}^{\infty} \lambda(A_j)$ for $A_j \in \mathcal{L}(\mathbb{R})$ 
 $A_j \cap A_i = \emptyset$ for $i \neq j$
- $\mathcal{L}(\mathbb{R})$ is larger than the Borel σ -algebra.
- If $A \in \mathcal{L}(\mathbb{R})$ with $\lambda(A) = 0$, (A is called null set)
then each $B \subseteq A$ satisfies $B \in \mathcal{L}(\mathbb{R})$.
- $\lambda([a, b]) = b - a$, $b \geq a$
- $\lambda(x + A) = \lambda(A)$ for all $x \in \mathbb{R}$, $A \in \mathcal{L}(\mathbb{R})$
(translation-invariant)

Definition (Lebesgue integral):


$$\int_A f d\lambda = \int_A f(x) d\lambda(x) = \int_A f(x) dx$$

defined by approximation with simple functions:







Multidimensional Integration - Part 2



Now go to \mathbb{R}^2 : $\mathbb{R} \times \mathbb{R}$ (construction of product measure)



- We get:
- product σ-algebra $\mathcal{L}(\mathbb{R}^2)$ (Lebesgue-measurable subsets of \mathbb{R}^2)
 - product measure $\lambda^{(2)}: \mathcal{L}(\mathbb{R}^2) \longrightarrow [0, \infty]$ Lebesgue measure on \mathbb{R}^2
 - $\lambda^{(2)}(A_1 \times A_2) = \lambda(A_1) \cdot \lambda(A_2)$ for $A_1 \in \mathcal{L}(\mathbb{R})$, $A_2 \in \mathcal{L}(\mathbb{R})$
 - properties like for the one-dimensional Lebesgue measure:
 - $\lambda^{(2)}(\emptyset) = 0$
 - $\lambda^{(2)}\left(\bigcup_{j=1}^{\infty} A_j\right) = \sum_{j=1}^{\infty} \lambda^{(2)}(A_j)$ for $A_j \in \mathcal{L}(\mathbb{R}^2)$, $A_j \cap A_i = \emptyset$ for $i \neq j$
 - $\mathcal{L}(\mathbb{R}^2)$ is larger than the Borel σ-algebra.
 - If $A \in \mathcal{L}(\mathbb{R}^2)$ with $\lambda^{(2)}(A) = 0$, (A is called null set)
then each $B \subseteq A$ satisfies $B \in \mathcal{L}(\mathbb{R}^2)$.
 - $\lambda^{(2)}([0,1] \times [0,1]) = 1$ (unit square as area 1)
 - $\lambda^{(2)}(x + A) = \lambda^{(2)}(A)$ for all $x \in \mathbb{R}^2$, $A \in \mathcal{L}(\mathbb{R}^2)$
(translation-invariant)




We call $\lambda^{(2)}$ the two-dimensional Lebesgue measure!

→ the corresponding Lebesgue integral: $\int_A f d\lambda^{(2)}$
the two-dimensional Lebesgue integral

Other notations:

$$\begin{aligned} \int_A f d\lambda^{(2)} &= \int_A f(x) d\lambda^{(2)}(x) \\ &= \int_A f(x_1, x_2) d\lambda^{(2)}(x_1, x_2) = \int_A f(x_1, x_2) d(x_1, x_2) \\ &= \int_A f(x) d^2x \end{aligned}$$

Do it again!



volume in \mathbb{R}^3 :


$\lambda^{(2)}(u) \cdot \lambda(A_3) \rightsquigarrow \lambda^{(3)}$ as a product measure

Result: n-dimensional Lebesgue measure on \mathbb{R}^n : $\lambda^{(n)}: \mathcal{L}(\mathbb{R}^n) \rightarrow [0, \infty]$

properties:

- $\lambda^{(n)}(\emptyset) = 0$
- $\lambda^{(n)}\left(\bigcup_{j=1}^{\infty} A_j\right) = \sum_{j=1}^{\infty} \lambda^{(n)}(A_j)$ for $A_j \in \mathcal{L}(\mathbb{R}^n)$, $A_j \cap A_i = \emptyset$ for $i \neq j$
- $\mathcal{L}(\mathbb{R}^n)$ is larger than the Borel σ -algebra.
- If $A \in \mathcal{L}(\mathbb{R}^n)$ with $\lambda^{(n)}(A) = 0$, (A is called null set)
then each $B \subseteq A$ satisfies $B \in \mathcal{L}(\mathbb{R}^n)$.
- $\lambda^{(n)}([0,1] \times [0,1] \times \dots \times [0,1]) = 1$
- $\lambda^{(n)}(x + A) = \lambda^{(n)}(A)$ for all $x \in \mathbb{R}^n$, $A \in \mathcal{L}(\mathbb{R}^n)$
(translation-invariant)


n-dimensional Lebesgue integral: $\int_A f d\lambda^{(n)} = \int_A f(x) d^n x$



Multidimensional Integration - Part 3

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}$$

Fubini's theorem



two-dimensional
Lebesgue integral

$$\int_U f(x_1, x_2) d(x_1, x_2)$$

Example:

$$\int_{[0,1] \times [0,2]} x_1 \cdot x_2 d(x_1, x_2) = \int_0^1 \left(\int_0^2 x_1 \cdot x_2 d x_2 \right) d x_1$$

two-dimensional
Lebesgue integral

one-dimensional
Lebesgue integral

one-dimensional
Lebesgue integral


Fubini's theorem (Fubini-Tonelli theorem)

Let $\lambda^{(n)}$ be the n -dimensional Lebesgue measure on \mathbb{R}^n

and $\lambda^{(m)}$ be the m -dimensional Lebesgue measure on \mathbb{R}^m .

Let $A \subseteq \mathbb{R}^n$, $B \subseteq \mathbb{R}^m$, and f be a measurable function with


either $f: A \times B \rightarrow [0, \infty]$



or $f: A \times B \rightarrow \mathbb{R}$ with $\int_{A \times B} |f| d\lambda^{(n+m)} < \infty$.

Then:

$$\int_{A \times B} f d\lambda^{(n+m)} = \int_A \left(\int_B f(x, y) d^m y \right) d^n x = \int_B \left(\int_A f(x, y) d^n x \right) d^m y$$



Multidimensional Integration - Part 4


Fubini's theorem (Fubini-Tonelli theorem): Let f be measurable with

either $f: A \times B \rightarrow [0, \infty]$ $(A \subseteq \mathbb{R}^n, B \subseteq \mathbb{R}^m)$
 or $f: A \times B \rightarrow \mathbb{R}$ with $\int_{A \times B} |f| d\lambda^{(n+m)} < \infty$.


Then:

$$\int_{A \times B} f d\lambda^{(n+m)} = \int_A \left(\int_B f(x, y) d^m y \right) d^n x = \int_B \left(\int_A f(x, y) d^n x \right) d^m y$$

Problem:



Example:




$$\int_U 1 \, d(x, y)$$

$$\underset{[0,1] \times [1,3]}{\int \tilde{f}(x, y) \, d(x, y)} \quad \text{with} \quad \tilde{f}(x, y) := \begin{cases} 1 & , x \in [0,1], y \in [1, 3-x^2] \\ 0 & \text{else} \end{cases}$$

$$\stackrel{\text{Fubini}}{=} \int_0^1 \left(\int_1^{3-x^2} \tilde{f}(x, y) \, dy \right) dx = \int_0^1 \left(\int_1^{3-x^2} 1 \, dy \right) dx$$


$$= \int_0^1 (3 - x^2 - 1) dx = \int_0^1 (2 - x^2) dx = \frac{5}{3}$$



Multidimensional Integration - Part 5

$f: U \rightarrow \mathbb{R}$ measurable, $U \subseteq \mathbb{R}^n$ open

$$\int_U f(x) d^n x$$



$\Phi: \tilde{U} \rightarrow U$ continuously differentiable + bijective

$\Phi^{-1}: U \rightarrow \tilde{U}$ continuously differentiable

substitution: $x = \Phi(\tilde{x}) \xrightarrow{\text{in one dimension!}} dx = \Phi'(x) d\tilde{x}$

$$\xrightarrow{\text{now!}} d^n x = |\det(J_{\Phi}(x))| d^n \tilde{x}$$

Change of variables formula: $\int_U f(x) d^n x = \int_{\tilde{U}} f(\Phi(\tilde{x})) |\det(J_{\Phi}(\tilde{x}))| d^n \tilde{x}$

If one exists,
then also the other!