


## Real Analysis - Part 10

### Bolzano-Weierstrass theorem

$(a_n)_{n \in \mathbb{N}}$  bounded  $\implies (a_n)_{n \in \mathbb{N}}$  has an accumulation value  
(has a convergent subsequence)




Proof:



If infinitely many sequence members in it: Choose left-hand interval  
Otherwise: Choose right-hand interval

New interval:



repeat

We get:  $[c_0, d_0] \supset [c_1, d_1] \supset [c_2, d_2] \supset [c_3, d_3] \supset \dots$

And:  $d_1 - c_1 = \frac{1}{2}(d_0 - c_0)$ ,  $d_2 - c_2 = \frac{1}{2}(d_1 - c_1) = \frac{1}{4}(d_0 - c_0)$ , ...

$$d_n - c_n = \frac{1}{2^n}(d_0 - c_0) \xrightarrow{n \rightarrow \infty} 0$$

We know:  $(c_n)_{n \in \mathbb{N}}$  mon. increasing and bounded  $\left. \begin{array}{l} (d_n)_{n \in \mathbb{N}} \text{ mon. decreasing and bounded} \end{array} \right\} \implies (c_n)_{n \in \mathbb{N}}, (d_n)_{n \in \mathbb{N}}$  convergent

By limit theorems:  $0 = \lim_{n \rightarrow \infty} (d_n - c_n) = \lim_{n \rightarrow \infty} d_n - \lim_{n \rightarrow \infty} c_n$

Define a subsequence  $(a_{n_k})_{k \in \mathbb{N}}$  by choosing  $a_{n_k} \in [c_k, d_k]$

$$\implies c_k \leq a_{n_k} \leq d_k$$

Sandwich theorem

$\implies (a_{n_k})_{k \in \mathbb{N}}$  is convergent