


Real Analysis - Part 10

Bolzano-Weierstrass theorem

$(a_n)_{n \in \mathbb{N}}$ bounded $\Rightarrow (a_n)_{n \in \mathbb{N}}$ has an accumulation value
(has a convergent subsequence)




Proof:



If infinitely many sequence members in it: Choose left-hand interval
Otherwise: Choose right-hand interval

New interval: $[c_1, d_1]$



We get: $[c_0, d_0] \supset [c_1, d_1] \supset [c_2, d_2] \supset [c_3, d_3] \supset \dots$

And: $d_1 - c_1 = \frac{1}{2}(d_0 - c_0)$, $d_2 - c_2 = \frac{1}{2}(d_1 - c_1) = \frac{1}{4}(d_0 - c_0)$, ...

$$d_n - c_n = \frac{1}{2^n}(d_0 - c_0) \xrightarrow{n \rightarrow \infty} 0$$

We know: $(c_n)_{n \in \mathbb{N}}$ mon. increasing and bounded } $\Rightarrow (c_n)_{n \in \mathbb{N}}, (d_n)_{n \in \mathbb{N}}$ convergent
 $(d_n)_{n \in \mathbb{N}}$ mon. decreasing and bounded

By limit theorems: $0 = \lim_{n \rightarrow \infty} (d_n - c_n) = \lim_{n \rightarrow \infty} d_n - \lim_{n \rightarrow \infty} c_n$

Define a subsequence $(a_{n_k})_{k \in \mathbb{N}}$ by choosing $a_{n_k} \in [c_k, d_k]$

$$\Rightarrow c_k \leq a_{n_k} \leq d_k$$

Sandwich theorem

$\Rightarrow (a_{n_k})_{k \in \mathbb{N}}$ is convergent