



The Bright Side of Mathematics

The following pages cover the whole Unbounded Operators course of the Bright Side of Mathematics. Please note that the creator lives from generous supporters and would be very happy about a donation. See more here: <https://tbsom.de/support>

Have fun learning mathematics!



Unbounded Operators - Part 1



- Motivation:
- partial differential equations
 - quantum mechanics: one needs operators X, P with

$$XP - PX = i \cdot I$$

Definition: Let $(X, \|\cdot\|_X)$, $(Y, \|\cdot\|_Y)$ be normed spaces (same field $F \in \{\mathbb{R}, \mathbb{C}\}$) and $\mathcal{D} \subseteq X$ subspace.

A linear map $T: \mathcal{D} \rightarrow Y$ is called an operator.

- Other notations:
- $T: X \supseteq \mathcal{D} \rightarrow Y$
 - $T: X \rightarrow Y$ with domain \mathcal{D}
 - (T, \mathcal{D}) or T with $\mathcal{D}(T) = \mathcal{D}$

Moreover: T is called densely defined if $\overline{\mathcal{D}}^{\|\cdot\|_X} = X$.

$$\text{Ran}(T) := \{Tx \mid x \in \mathcal{D}\} \subseteq Y \text{ subspace}$$

$$\text{Ker}(T) := \{x \in \mathcal{D} \mid Tx = 0\} \subseteq X \text{ subspace}$$

T is called bounded if $\exists C > 0 \forall x \in \mathcal{D} : \|Tx\|_Y \leq C \cdot \|x\|_X$

T is called unbounded if $\forall C > 0 \exists x \in \mathcal{D} : \|Tx\|_Y > C \cdot \|x\|_X$

Recall: T is bounded $\Leftrightarrow T$ is continuous at all points $x \in \mathcal{D}$

Therefore: T is unbounded $\Leftrightarrow T$ is not continuous (at no point $x \in \mathcal{D}$)



Unbounded Operators - Part 2

Recall: operator $T: X \rightarrow Y$ with $\mathcal{D}(T) = \mathcal{D}$

means: $T: \mathcal{D} \rightarrow Y$ linear map


Fact: If $\text{Ker}(T) = \{0\}$, then $T^{-1}: Y \rightarrow X$ with $\mathcal{D}(T^{-1}) = \text{Ran}(T)$
 \hookrightarrow always defined as an operator

Examples: $X = Y = C([0,1])$ (with supremum norm $\|\cdot\|_\infty$)

(a) $T: X \rightarrow Y$ with $\mathcal{D}(T) = C^1([0,1])$

$$Tx = x'$$

unbounded operator



$$\|T\| = \sup_{\|x\|_\infty=1} \|Tx\|_\infty = \sup_{\|x\|_\infty=1} \|x'\|_\infty = \infty$$

(b) $S: X \rightarrow Y$ with $\mathcal{D}(S) = \{x \in C^1([0,1]) \mid x(0) = 0\}$

$$Sx = x'$$

notations: $S \subseteq T$

the operator T is an extension of S
 the operator S is a restriction of T

Note: • $\text{Ker}(T) \neq \{0\}$ not injective:

• $\text{Ker}(S) = \{0\}$ injective! $\Rightarrow S^{-1}$ exists

• T is densely defined $\left(\overline{C^1([0,1])}^{\|\cdot\|_\infty} = C([0,1]) \right)$


• S is not densely defined



Unbounded Operators - Part 3

Recall: operator $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ (linear map between normed spaces)

\Leftrightarrow subset in $X \times Y$



graph of T : $G_T := \{(x, y) \in X \times Y \mid x \in \mathcal{D}(T), Tx = y\}$

$X \times Y$ normed space with $\|(x, y)\|_{X \times Y} := \|x\|_X + \|y\|_Y$

Definition: An operator $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ is called a closed operator if the graph G_T is closed (in the normed space $X \times Y$).

Note: T closed \Leftrightarrow


for each sequence $(x_n) \subseteq \mathcal{D}(T)$ with
 $x_n \xrightarrow{n \rightarrow \infty} x \in X$, $Tx_n \rightarrow y \in Y$,
 we have: $x \in \mathcal{D}(T)$ and $Tx = y$

Proof: G_T closed \Leftrightarrow for each sequence $(x_n, Tx_n) \subseteq G_T$
 that is convergent in $X \times Y$ with limit

$(x, y) \in X \times Y$,

we have: $(x, y) \in G_T$.

$x \in \mathcal{D}(T)$ and $Tx = y$



Remember: $T: X \rightarrow Y$ with $\mathcal{D}(T) = X$ bounded \Rightarrow closed operator



Unbounded Operators - Part 4

Closed operator: $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ closed

$$\Leftrightarrow G_T := \{(x, y) \in X \times Y \mid x \in \mathcal{D}(T), Tx = y\} \text{ closed}$$

Closable operator: $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ closable

$$:\Leftrightarrow \overline{G_T} \text{ is the graph of an operator } \overline{T} \leftarrow \text{closure of } T$$

Proposition: $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ closable

$$\Leftrightarrow \overline{G_T} \text{ is a graph (not possible } (0, 0), (0, y) \in \overline{G_T} \text{ for } y \neq 0)$$

$$\Leftrightarrow \text{If } (0, y) \in \overline{G_T}, \text{ then } y = 0. \quad \boxed{G_T := \{(x, y) \in X \times Y \mid x \in \mathcal{D}(T), Tx = y\}}$$

$$\Leftrightarrow \text{For each } (x_n) \subseteq \mathcal{D}(T) \text{ with } x_n \rightarrow 0 \text{ and } Tx_n \rightarrow y, \\ \text{we have } y = 0.$$

Define \overline{T} for a closable operator $T: X \supseteq \mathcal{D}(T) \rightarrow Y$:

$$\mathcal{D}(\overline{T}) := \{x \in X \mid \exists (x_n) \subseteq \mathcal{D}(T) : x_n \rightarrow x \text{ and } Tx_n \text{ convergent}\}$$

$$\overline{T}x := \lim_{n \rightarrow \infty} Tx_n \quad \text{operator! (closure of } T)$$

$$\Rightarrow T \subseteq \overline{T}$$



Unbounded Operators - Part 6

Closed Graph Theorem: X, Y Banach spaces, $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ operator with $\mathcal{D}(T)$ closed (e.g. $\mathcal{D}(T) = X$).

Then: T closed $\Leftrightarrow T$ continuous (bounded)

Proof: Assume: $\mathcal{D}(T) = X$.

(\Leftarrow) Choose $(x_n) \subseteq \mathcal{D}(T)$ with $x_n \rightarrow x \in X$ and $Tx_n \rightarrow y \in Y$

$$\stackrel{T \text{ continuous}}{\Rightarrow} y = \lim_{n \rightarrow \infty} T(x_n) = T(\lim_{n \rightarrow \infty} x_n) = Tx$$


$$\Rightarrow x \in \mathcal{D}(T) \text{ and } Tx = y \Rightarrow T \text{ closed}$$

(\Rightarrow) Assume T is closed $\Rightarrow G_T$ is closed in $X \times Y \Rightarrow (G_T, \|\cdot\|_{X \times Y})$ Banach space


Define operators: $P_X: G_T \rightarrow X$ and $P_Y: G_T \rightarrow Y$ linear + bounded
 $(x, y) \mapsto x$ $(x, y) \mapsto y$
 bijective!

Bounded
Inverse
Theorem

Functional Analysis
- Part 27



$P_X^{-1}: X \rightarrow G_T$ is continuous (bounded operator)
 $x \mapsto (x, Tx)$



$T = P_Y P_X^{-1}$ composition of continuous maps

$\Rightarrow T$ continuous (bounded)



Unbounded Operators - Part 7

X, Y Banach spaces, $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ operator

Closed Graph Theorem: $\mathcal{D}(T) = X \implies (T \text{ closed} \iff T \text{ bounded})$

Example: functional $T: X \rightarrow \mathbb{C}$ unbounded (see part 5)

\hookrightarrow extend: $\mathcal{D}(T) = X$

$\implies T$ not closed

Proposition: X, Y Banach spaces, $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ operator.

Then: T closed $\iff (\mathcal{D}(T), \|\cdot\|_T)$ complete

\uparrow graph norm

$$\|x\|_T := \|x\|_X + \|Tx\|_Y$$

Proof: $J: (\mathcal{D}(T), \|\cdot\|_T) \longrightarrow (G_T, \|\cdot\|_{X \times Y})$ } linear + bijective

$$x \longmapsto (x, Tx)$$

$$\|Jx\|_{X \times Y} = \|(x, Tx)\|_{X \times Y} = \|x\|_X + \|Tx\|_Y = \|x\|_T$$

$\implies J$ is an isometric isomorphism

$(\mathcal{D}(T), \|\cdot\|_T)$ complete $\iff (G_T, \|\cdot\|_{X \times Y})$ complete

$\iff (G_T, \|\cdot\|_{X \times Y})$ closed in $X \times Y$

$\iff T$ closed



Unbounded Operators - Part 8

For bounded operators: $T: X \rightarrow Y \rightsquigarrow T^*: Y \rightarrow X$ adjoint
Hilbert spaces $\langle y, Tx \rangle_Y = \langle T^*y, x \rangle_X$

$T: X \rightarrow Y \rightsquigarrow T': Y' \rightarrow X'$ adjoint
Banach spaces $T'(y')(x) = y'(Tx)$
 for $y' \in Y', x \in X$

Proposition: X, Y Banach spaces, $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ densely defined operator
 $\hookrightarrow \overline{\mathcal{D}(T)} = X$

Then there is an operator $T': Y' \supseteq \mathcal{D}(T') \rightarrow X'$ with

$$y'(Tx) = T'(y')(x) \text{ for } x \in \mathcal{D}(T), y' \in \mathcal{D}(T').$$

The domain $\mathcal{D}(T')$ can be chosen maximally.

Proof: set $\mathcal{D}(T') := \{y' \in Y' \mid \text{there is } x' \in X' \text{ with } y'(Tx) = x'(x) \text{ for all } x \in \mathcal{D}(T)\}$

and define: $T'(y') := x'$

Well-defined? Assume there are $x'_1, x'_2 \in X'$ with $y'(Tx) = x'_1(x)$ for all $x \in \mathcal{D}(T)$
 $y'(Tx) = x'_2(x)$

$$\Rightarrow x'_1(x) = x'_2(x) \text{ for all } x \in \mathcal{D}(T)$$

$$\Rightarrow (x'_1 - x'_2)(x) = 0 \text{ for all } x \in \mathcal{D}(T) \xrightarrow[\text{continuity}]{\text{dense}} (x'_1 - x'_2)(x) = 0 \text{ for all } x \in X$$

$$\Rightarrow x'_1 = x'_2 \quad \square$$

For Hilbert spaces: X, Y Hilbert spaces, $T: X \supseteq \mathcal{D}(T) \rightarrow Y$ densely defined operator
 $\hookrightarrow \overline{\mathcal{D}(T)} = X$

$$\mathcal{D}(T^*) := \{y \in Y \mid \text{there is } \tilde{x} \in X \text{ with } \langle y, Tx \rangle_Y = \langle \tilde{x}, x \rangle_X \text{ for all } x \in \mathcal{D}(T)\}$$

$$T^*(y) := \tilde{x}$$