

Unbounded Operators - Part 2

Recall: operator $T: X \rightarrow Y$ with $\mathcal{D}(T) = \mathcal{D}$

means: $T: \mathcal{D} \rightarrow Y$ linear map


Fact: If $\text{Ker}(T) = \{0\}$, then $T^{-1}: Y \rightarrow X$ with $\mathcal{D}(T^{-1}) = \text{Ran}(T)$
↳ always defined as an operator

Examples: $X = Y = C([0,1])$ (with supremum norm $\|\cdot\|_\infty$)

(a) $T: X \rightarrow Y$ with $\mathcal{D}(T) = C^1([0,1])$

$$Tx = x'$$

unbounded operator



$$\|T\| = \sup_{\|x\|_\infty=1} \|Tx\|_\infty = \sup_{\|x\|_\infty=1} \|x'\|_\infty = \infty$$

(b) $S: X \rightarrow Y$ with $\mathcal{D}(S) = \{x \in C^1([0,1]) \mid x(0) = 0\}$

$$Sx = x'$$

notations: $S \subseteq T$

the operator T is an extension of S
the operator S is a restriction of T

Note: • $\text{Ker}(T) \neq \{0\}$ not injective!

• $\text{Ker}(S) = \{0\}$ injective! $\Rightarrow S^{-1}$ exists

• T is densely defined $\left(\overline{C^1([0,1])}^{\|\cdot\|_\infty} = C([0,1]) \right)$

• S is not densely defined