




The Bright Side of Mathematics

Probability Theory - Part 8

Bayes's theorem: $P(A|B) = \frac{P(A \cap B)}{P(B)}$, $P(B|A) = \frac{P(B \cap A)}{P(A)}$


$$\Rightarrow P(A|B) \cdot P(B) = P(B|A) \cdot P(A)$$

Law of total probability: (Ω, \mathcal{A}, P) probability space




$$P(A) = P(\underbrace{(A \cap B) \cup (A \cap B^c)}_{\text{disjoint union}}) = P(A \cap B) + P(A \cap B^c) = P(A|B) \cdot P(B) + P(A|B^c) \cdot P(B^c)$$

Case with countably many sets: $B_i \in \mathcal{A}$ for $i \in I \subseteq \mathbb{N}$ with $\bigcup_{i \in I} B_i = \Omega$
disjoint union



$$P(A) = P\left(\bigcup_{i \in I} (A \cap B_i)\right) = \sum_{i \in I} P(A \cap B_i) = \sum_{i \in I} P(A|B_i) \cdot P(B_i)$$

Example: Monty Hall problem



Second: Show master opens a door with a goat (3) (never the door you picked)

Third: Stay or switch

$C_j :=$ car is behind door j , $S_j :=$ show master opens door j (in the second step)

We know: $P(S_3|C_3) = 0$, $P(S_3|C_2) = 1$, $P(S_3|C_1) = \frac{1}{2}$

$$P(C_2|S_3) \stackrel{\substack{\uparrow \\ \text{Bayes's} \\ \text{theorem}}}{=} \frac{P(S_3|C_2) \cdot P(C_2)}{P(S_3)} \stackrel{\substack{\uparrow \\ \text{Law of total} \\ \text{probability}}}{=} \frac{P(S_3|C_2) \cdot P(C_2)}{\sum_{j=1}^3 P(S_3|C_j) \cdot P(C_j)}$$

$$= \frac{P(S_3|C_2) \cdot P(C_2)}{P(S_3|C_1) \cdot P(C_1) + P(S_3|C_2) \cdot P(C_2) + P(S_3|C_3) \cdot P(C_3)} = \frac{2}{3}$$

Annotations in the diagram:
- $P(S_3|C_1) = \frac{1}{2}$
- $P(C_1) = \frac{1}{3}$
- $P(S_3|C_2) = 1$
- $P(C_2) = \frac{1}{3}$
- $P(S_3|C_3) = 0$
- $P(C_3) = \frac{1}{3}$