



The Bright Side of Mathematics

Real Analysis - Part 57

Integration by substitution

$I \subseteq \mathbb{R}$ interval, $f: I \rightarrow \mathbb{R}$ continuous, $\phi: [a, b] \rightarrow I$ continuously differentiable

Then:
$$\int_a^b f(\phi(t)) \cdot \phi'(t) dt = \int_{\phi(a)}^{\phi(b)} f(x) dx$$

Remember: $x = \phi(t)$

$$\frac{dx}{dt} = \phi'(t) \Rightarrow dx = \phi'(t) dt$$

Example:
$$\int_0^1 t^2 \cdot \sin(t^3) dt = \frac{1}{3} \int_0^1 \sin(t^3) \cdot 3t^2 dt = \frac{1}{3} \int_0^1 \sin(x) dx$$

$x = t^3$
 $dx = 3t^2 dt$

Proof: Let $F: I \rightarrow \mathbb{R}$ be an antiderivative of f

$$(F \circ \phi)'(t) \stackrel{\text{chain rule}}{=} F'(\phi(t)) \cdot \phi'(t) = f(\phi(t)) \cdot \phi'(t)$$

$$\int_a^b f(\phi(t)) \cdot \phi'(t) dt = \int_a^b (F \circ \phi)'(t) dt = (F \circ \phi)(t) \Big|_{t=a}^{t=b}$$

$$= F(x) \Big|_{x=\phi(a)}^{x=\phi(b)} = \int_{\phi(a)}^{\phi(b)} f(x) dx \quad \square$$

Another substitution rule: $f: [a, b] \rightarrow \mathbb{R}$ continuous, $\phi: J \rightarrow I$ continuously differentiable and bijective
 $J, I \subseteq \mathbb{R}$ intervals, $I \supseteq [a, b]$

$$\int_a^b f(x) dx = \int_{\phi^{-1}(a)}^{\phi^{-1}(b)} f(\phi(t)) \cdot \phi'(t) dt$$

Example:

$b \in [0, 1]$

$$\int_0^b \frac{1}{\sqrt{1-x^2}} dx$$

try: $\phi(t) = \sin(t)$

substitution: $x = \sin(t)$

$$dx = \cos(t) dt$$

$$\sin^2(t) + \cos^2(t) = 1$$

$$= \int_0^{\arcsin(b)} \frac{1}{\underbrace{\sqrt{1-\sin^2(t)}}_{\cos(t)}} \cos(t) dt = \int_0^{\arcsin(b)} 1 dt = \arcsin(b)$$

bijective: $[0, \frac{\pi}{2}] \rightarrow [0, 1]$